

ETAR (F-12): sc-515948

BACKGROUND

Endothelin receptor A (ETAR), also known as EDNRA, ET1 receptor, ETA, EDN1 and ET-AR, is a member of the guanine-binding regulatory protein-coupled receptor family. ETAR binds endothelins and has the highest affinity for its ligand, ET1, as compared to the ETBR receptor. Both ET receptors, ETAR and ETBR, are activated by ET1, which results in inhibition of active lens sodium-potassium transport. Activation of the ET receptors also causes an increase in cytoplasmic calcium concentration in cultured lens epithelial cells. In addition, ETAR induces arachidonic acid accumulation. ETAR has seven hydrophobic transmembrane domains and is expressed in aorta, lung, atrium, kidney, placenta and prostate. Specifically, placental vascular smooth muscle cells (PVSMCs) exclusively express ETAR.

REFERENCES

1. Adachi, M., et al. 1991. Cloning and characterization of cDNA encoding human A-type endothelin receptor. *Biochem. Biophys. Res. Commun.* 180: 1265-1272.
2. Lin, H., et al. 1991. Cloning and functional expression of a vascular smooth muscle endothelin 1 receptor. *Proc. Natl. Acad. Sci. USA* 88: 3185-3189.
3. Kobayashi, S., et al. 1994. Binding and functional properties of endothelin receptor subtypes in the human prostate. *Mol. Pharmacol.* 45: 306-311.
4. Miyamoto, Y., et al. 1996. Alternative RNA splicing of the human endothelin-A receptor generates multiple transcripts. *Biochem. J.* 313: 795-801.
5. Okafor, M., et al. 2001. The inhibitory influence of endothelin on active sodium-potassium transport in porcine lens. *Invest. Ophthalmol. Vis. Sci.* 42: 1018-1023.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 131243. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: *Ednra* (mouse) mapping to 8 C1.

SOURCE

ETAR (F-12) is a mouse monoclonal antibody raised against amino acids 21-80 mapping within an N-terminal extracellular domain of ETAR of mouse origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ETAR (F-12) is available conjugated to agarose (sc-515948 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-515948 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515948 PE), fluorescein (sc-515948 FITC), Alexa Fluor® 488 (sc-515948 AF488), Alexa Fluor® 546 (sc-515948 AF546), Alexa Fluor® 594 (sc-515948 AF594) or Alexa Fluor® 647 (sc-515948 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-515948 AF680) or Alexa Fluor® 790 (sc-515948 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

ETAR (F-12) is recommended for detection of ETAR of mouse and rat origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ETAR siRNA (m): sc-39961, ETAR siRNA (r): sc-270097, ETAR shRNA Plasmid (m): sc-39961-SH, ETAR shRNA Plasmid (r): sc-270097-SH, ETAR shRNA (m) Lentiviral Particles: sc-39961-V and ETAR shRNA (r) Lentiviral Particles: sc-270097-V.


Molecular Weight of ETAR: 69 kDa.

Positive Controls: ETAR (m): CHO Lysate: sc-110170.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ETAR (F-12): sc-515948. Western blot analysis of ETAR expression in non-transfected: sc-117750 (A) and mouse ETAR transfected: sc-110170 (B) CHO whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA