

# p87PIKAP (1G16): sc-517014

## BACKGROUND

p87PIKAP (p87 PI3K adapter protein), also known as PIK3R6 (phosphoinositide-3-kinase, regulatory subunit 6), p84 or HsT41028, is a 754 amino acid cytoplasmic protein that functions as a regulatory subunit of the PI 3-kinase p110 $\gamma$  complex. Expressed in heart, dendritic cells, macrophages and neutrophils, p87PIKAP drives PI 3-kinase p110 $\gamma$  activation, interacts with PDE3B and is thought to cause the two proteins to form a scaffolding interaction. The gene encoding p87PIKAP maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

## REFERENCES

- Hall, J.M., Friedman, L., Guenther, C., Lee, M.K., Weber, J.L., Black, D.M. and King, M.C. 1992. Closing in on a breast cancer gene on chromosome 17q. *Am. J. Hum. Genet.* 50: 1235-1242.
- Stoyanov, B., Volinia, S., Hanck, T., Rubio, I., Loubtchenkov, M., Malek, D., Stoyanova, S., Vanhaesebroeck, B., Dhand, R. and Nürnberg, B. 1995. Cloning and characterization of a G protein-activated human phosphoinositide-3 kinase. *Science* 269: 690-693.
- Kersemaekers, A.M., Hermans, J., Fleuren, G.J. and van de Vijver, M.J. 1998. Loss of heterozygosity for defined regions on chromosomes 3, 11 and 17 in carcinomas of the uterine cervix. *Br. J. Cancer* 77: 192-200.
- Soussi, T., Dehouche, K. and Beroud, C. 2000. p53 website and analysis of p53 gene mutations in human cancer: forging a link between epidemiology and carcinogenesis. *Hum. Mutat.* 15: 105-113.
- Piura, B., Rabinovich, A. and Yanai-Inbar, I. 2001. Three primary malignancies related to BRCA mutation successively occurring in a BRCA1 185delAG mutation carrier. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 97: 241-244.
- Minamoto, T., Buschmann, T., Habelhah, H., Matusevich, E., Tahara, H., Boerresen-Dale, A.L., Harris, C., Sidransky, D. and Ronai, Z. 2001. Distinct pattern of p53 phosphorylation in human tumors. *Oncogene* 20: 3341-3347.
- Brock, C., Schaefer, M., Reusch, H.P., Czupalla, C., Michalke, M., Spicher, K., Schultz, G. and Nürnberg, B. 2003. Roles of G $\beta$  $\gamma$  in membrane recruitment and activation of p110 $\gamma$ /p101 phosphoinositide 3-kinase  $\gamma$ . *J. Cell Biol.* 160: 89-99.
- Suire, S., Coadwell, J., Ferguson, G.J., Davidson, K., Hawkins, P. and Stephens, L. 2005. p84, a new G $\beta$  $\gamma$ -activated regulatory subunit of the type IB phosphoinositide 3-kinase p110 $\gamma$ . *Curr. Biol.* 15: 566-570.
- Voigt, P., Dörner, M.B. and Schaefer, M. 2006. Characterization of p87PIKAP, a novel regulatory subunit of phosphoinositide 3-kinase  $\gamma$  that is highly expressed in heart and interacts with PDE3B. *J. Biol. Chem.* 281: 9977-9986.

## CHROMOSOMAL LOCATION

Genetic locus: PIK3R6 (human) mapping to 17p13.1.

## SOURCE

p87PIKAP (1G16) is a mouse monoclonal antibody raised against amino acids 661-754 representing partial length p87PIKAP of human origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG $_1$  kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

p87PIKAP (1G16) is recommended for detection of p87PIKAP of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for p87PIKAP siRNA (h): sc-93748, p87PIKAP shRNA Plasmid (h): sc-93748-SH and p87PIKAP shRNA (h) Lentiviral Particles: sc-93748-V.

Molecular Weight of p87PIKAP: 84 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



p87PIKAP (1G16): sc-517014. Western blot analysis of human recombinant p87PIKAP fusion protein.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.