p35 (G-7): sc-518015



The Power to Question

BACKGROUND

Cyclin dependent kinase-5 (Cdk5), a key regulator of cell cycle progression, was originally isolated on the basis of its structural homology to Cdc2, a well-characterized regulator of cell cycle progression. Although Cdk5 is expressed at the highest level in the brain of adult mice, intermediate levels in testis and low or undetectable levels in all other tissues, brain is the only tissue from which Cdk5 can be isolated as an active kinase. These findings may be explained by the cloning and characterization of a Cdk5 regulatory subunit, designated p35. p35 displays a neuronal cell-specific pattern of expression, physically associates with Cdk5 and activates Cdk5 enzymatic activity. p35 is also expressed in many tissues in a truncated form, designated p25.

REFERENCES

- Murray, A.W. and Kirschner, M.W. 1989. Dominoes and clocks: the union of two views of the cell cycle. Science 246: 614-621.
- Nurse, P. 1990. Universal control mechanism regulating onset of M-phase. Nature 344: 503-508.
- 3. Pines, J. and Hunter, T. 1990. Cdc2 p34: the S and M kinase? New Biol. 2: 389-401
- 4. Draetta, G. 1990. Cell cycle control in eukaryotes: molecular mechanisms of Cdc2 activation. Trends Biochem. Sci. 15: 378-383.
- 5. Meyerson, M., Enders, G.H., Wu, C.L., Su, L.K., Gorka, C., Nelson, C., Harlow, E. and Tsai, L.H. 1992. A family of human Cdc2-related protein kinases. EMBO J. 11: 2909-2917.
- Tsai, L.H., Delalle, I., Caviness, V.S., Jr., Chae, T. and Harlow, E. 1994. p35 is a neural-specific regulatory subunit of cyclin-dependent kinase 5. Nature 371: 419-423.
- Lew, J., Huang, Q.Q., Qi, Z., Winkfein, R.J., Aebersold, R., Hunt, T. and Wang, J.H. 1994. A brain-specific activator of cyclin-dependent kinase 5. Nature 371: 423-426.

CHROMOSOMAL LOCATION

Genetic locus: CDK5R1 (human) mapping to 17q11.2.

SOURCE

p35 (G-7) is a mouse monoclonal antibody raised against amino acids 69-140 mapping near the N-terminus of p35 of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

p35 (G-7) is recommended for detection of p35 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for p35 siRNA (h): sc-36153, p35 shRNA Plasmid (h): sc-36153-SH and p35 shRNA (h) Lentiviral Particles: sc-36153-V.

Molecular Weight of p35 truncated form: 25 kDa.


Molecular Weight of full length p35 precursor: 35 kDa.


Positive Controls: mouse p35 transfected HEK293T whole cell lysate.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG κ BP-FITC: sc-516140 or m-lgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA





p35 (G-7): sc-518015. Western blot analysis of human recombinant p35 fusion protein.

p35 (G-7): sc-518015. Western blot analysis of p35 expression in non-transfected (**A**) and mouse p35 transfected (**B**) HEK293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.