

A cyclase III (E-8): sc-518057

BACKGROUND

Adenylyl cyclases function to convert ATP to cyclic AMP in response to activation by a variety of hormones, neurotransmitters and other regulatory molecules. Cyclic AMP, in turn, activates several other target molecules to control a broad range of diverse phenomena such as metabolism, gene transcription and memory. Adenylyl cyclases respond to receptor-initiated signals, mediated by the G_s and G_i heterotrimeric G proteins. The binding of an agonist to a G_s -coupled receptor catalyzes the exchange of GDP (bound to $G_{\alpha s}$) for GTP, the dissociation of $GTP-G_{\alpha s}$ from $G_{\beta\gamma}$ and $G_{\alpha s}$ -mediated activation of adenylyl cyclase. Adenylyl cyclases of the type II family differ from other subforms in that they are conditionally stimulated by $G_{\alpha s/\beta\gamma}$ subunits and regulated by PKC-mediated C-terminal phosphorylation. Both short- and long-term activation of D_{2L} dopamine receptors result in a marked degree of sensitization of A cyclase I, II, V and IX, but not A cyclase VIII. The effects on A cyclase I, II and VIII is dependent upon the ability of these A cyclase isoforms to synergistically respond to selective activators in the presence of activated $G_{\alpha s}$. Belonging to the adenylyl cyclase class IV family, A cyclase III is activated by G_{olf} , which results in an elevation of cyclic AMP and subsequent activation of a cyclic nucleotide-gated channel.

REFERENCES

1. Gilman, A.G. 1987. G proteins: transducers of receptor-generated signals. *Annu. Rev. Biochem.* 56: 615-649.
2. Bourne, H.R., et al. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
3. Tang, W.J., et al. 1992. Adenylyl cyclases. *Cell* 70: 869-872.
4. Taussig, R., et al. 1994. Distinct patterns of bidirectional regulation of mammalian adenylyl cyclases. *J. Biol. Chem.* 269: 6093-6100.
5. Liu, C.Y., et al. 1999. FICRHR/cyclic AMP signaling in myenteric ganglia and calbindin-D28 intrinsic primary afferent neurons involves adenylyl cyclases I, III and IV. *Brain Res.* 826: 253-269.
6. Gibson, A.D. and Garbers, D.L. 2000. Guanylyl cyclases as a family of putative odorant receptors. *Annu. Rev. Neurosci.* 23: 417-439.
7. Parkinson, N.A., et al. 2001. A nuclear location for Ca^{2+} -activated adenylyl cyclases I and III in neurons. *Brain Res. Mol. Brain Res.* 91: 43-49.
8. Cumbay, M.G., et al. 2001. Heterologous sensitization of recombinant adenylyl cyclases by activation of D_2 dopamine receptors. *J. Pharmacol. Exp. Ther.* 297: 1201-1209.

CHROMOSOMAL LOCATION

Genetic locus: ADCY3 (human) mapping to 2p23.3; Adcy3 (mouse) mapping to 12 A1.1.

SOURCE

A cyclase III (E-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 24-44 within an N-terminal cytoplasmic domain of A cyclase III of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

A cyclase III (E-8) is available conjugated to agarose (sc-518057 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-518057 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518057 PE), fluorescein (sc-518057 FITC), Alexa Fluor[®] 488 (sc-518057 AF488), Alexa Fluor[®] 546 (sc-518057 AF546), Alexa Fluor[®] 594 (sc-518057 AF594) or Alexa Fluor[®] 647 (sc-518057 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-518057 AF680) or Alexa Fluor[®] 790 (sc-518057 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS


A cyclase III (E-8) is recommended for detection of A cyclase III of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for A cyclase III siRNA (h): sc-29600, A cyclase III siRNA (m): sc-29601, A cyclase III shRNA Plasmid (h): sc-29600-SH, A cyclase III shRNA Plasmid (m): sc-29601-SH, A cyclase III shRNA (h) Lentiviral Particles: sc-29600-V and A cyclase III shRNA (m) Lentiviral Particles: sc-29601-V.


Molecular Weight of A cyclase III glycosylated forms: 170/180 kDa.

Positive Controls: U-87 MG cell lysate: sc-2411, HISM cell lysate: sc-2229 or WI-38 whole cell lysate: sc-364260.

DATA



A cyclase III (E-8): sc-518057. Western blot analysis of A cyclase III expression in HISM (A), U-87 MG (B) and WI-38 (C) whole cell lysates.



A cyclase III (E-8): sc-518057. Western blot analysis of A cyclase III expression in non-transfected (A) and mouse A cyclase III transfected (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.