

# MB67 (C-12): sc-518072

## BACKGROUND

The CAR "constitutively acting receptor" proteins, CAR1 and CAR2, are mouse nuclear hormone receptors. CAR1 and CAR2, along with their human homolog, MB67, are in highest expression in the liver and belong to a group of receptors known as orphan receptors due to their lack of a known ligand. Unlike conventional hormone receptors which activate transcription upon binding with steroids, retinoids, and thyroid hormones, the CAR and MB67 orphan receptors are transcriptionally active in the absence of exogenous hormone. The CAR and MB67 orphan receptors bind to DNA in the form of a heterodimer with the retinoic-X receptor and activate gene transcription in a constitutive manner.

## REFERENCES

1. Davies, P. and Rushmere, N.K. 1988. The structure and function of steroid receptors. *Sci. Prog.* 72: 563-578.
2. Baes, M., Gulick, T., Choi, H.S., Martinoli, M.G., Simha, D. and Moore, D.D. 1994. A new orphan member of the nuclear hormone receptor superfamily that interacts with a subset of retinoic acid response elements. *Mol. Cell. Biol.* 14: 1544-1551.
3. Mangelsdorf, D.J. and Evans, R.M. 1995. The RXR heterodimers and orphan receptors. *Cell* 83: 841-850.
4. Choi, H.S., Chung, M., Tzameli, I., Simha, D., Lee, Y.K., Seol, W. and Moore, D.D. 1997. Differential transactivation by two isoforms of the orphan nuclear hormone receptor CAR. *J. Biol. Chem.* 272: 23565-23571.
5. Forman, B.M., Tzameli, I., Choi, H.S., Chen, J., Simha, D., Seol, W., Evans, R.M. and Moore, D.D. 1998. Androstane metabolites bind to and deactivate the nuclear receptor CAR- $\beta$ . *Nature* 395: 612-615.

## CHROMOSOMAL LOCATION

Genetic locus: NR1I3 (human) mapping to 1q23.3.

## SOURCE

MB67 (C-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 1-24 at the N-terminus of MB67 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>3</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

MB67 (C-12) is available conjugated to agarose (sc-518072 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-518072 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518072 PE), fluorescein (sc-518072 FITC), Alexa Fluor<sup>®</sup> 488 (sc-518072 AF488), Alexa Fluor<sup>®</sup> 546 (sc-518072 AF546), Alexa Fluor<sup>®</sup> 594 (sc-518072 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-518072 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-518072 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-518072 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS


MB67 (C-12) is recommended for detection of MB67 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MB67 siRNA (h): sc-39918, MB67 shRNA Plasmid (h): sc-39918-SH and MB67 shRNA (h) Lentiviral Particles: sc-39918-V.


Molecular Weight of MB67: 40 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

## DATA



MB67 (C-12): sc-518072. Western blot analysis of MB67 expression in HeLa whole cell lysate.



MB67 (C-12): sc-518072. Western blot analysis of human recombinant MB67.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.