

$G_{\alpha 13}$ (A-6): sc-518113

BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (i.e., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein α , β and γ polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their α subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Four distinct classes of G_{α} subunits have been identified; these include $G_{\alpha s}$, $G_{\alpha i}$, $G_{\alpha q}$ and $G_{\alpha 12/13}$. The two members of the fourth class of G_{α} subunit proteins, $G_{\alpha 12}$ and $G_{\alpha 13}$, are insensitive to ADP-ribosylation by pertussis toxin, share 67% identity with each other and less than 45% identity with other G_{α} subunits and are widely expressed in a broad range of tissues.

REFERENCES

1. Strathmann, M., et al. 1989. Diversity of the G-protein family: sequences from five additional α subunits in the mouse. *Proc. Natl. Acad. Sci. USA* 86: 7407-7409.
2. Strathmann, M.P. and Simon, M.I. 1991. $G_{\alpha 12}$ and $G_{\alpha 13}$ subunits define a fourth class of G protein α subunits. *Proc. Natl. Acad. Sci. USA* 88: 5582-5586.
3. Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
4. von Weizsäcker, E., et al. 1992. Diversity among the β subunits of heterotrimeric GTP-binding proteins: characterization of a novel β -subunit cDNA. *Biochem. Biophys. Res. Commun.* 183: 350-356.

CHROMOSOMAL LOCATION

Genetic locus: GNA13 (human) mapping to 17q24.1; Gna13 (mouse) mapping to 11 E1.

SOURCE

$G_{\alpha 13}$ (A-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 194-216 within an internal region of $G_{\alpha 13}$ of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

$G_{\alpha 13}$ (A-6) is available conjugated to agarose (sc-518113 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-518113 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518113 PE), fluorescein (sc-518113 FITC), Alexa Fluor® 488 (sc-518113 AF488), Alexa Fluor® 546 (sc-518113 AF546), Alexa Fluor® 594 (sc-518113 AF594) or Alexa Fluor® 647 (sc-518113 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-518113 AF680) or Alexa Fluor® 790 (sc-518113 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

$G_{\alpha 13}$ (A-6) is recommended for detection of $G_{\alpha 13}$ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for $G_{\alpha 13}$ siRNA (h): sc-35427, $G_{\alpha 13}$ siRNA (m): sc-35428, $G_{\alpha 13}$ shRNA Plasmid (h): sc-35427-SH, $G_{\alpha 13}$ shRNA Plasmid (m): sc-35428-SH, $G_{\alpha 13}$ shRNA (h) Lentiviral Particles: sc-35427-V and $G_{\alpha 13}$ shRNA (m) Lentiviral Particles: sc-35428-V.

Molecular Weight of $G_{\alpha 13}$: 44 kDa.


Positive Controls: $G_{\alpha 13}$ (h): 293T Lysate: sc-115367, $G_{\alpha 13}$ (m): 293T Lysate: sc-115358 or HEK293T whole cell lysate: sc-45137.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



$G_{\alpha 13}$ (A-6): sc-518113. Western blot analysis of $G_{\alpha 13}$ expression in non-transfected: sc-117752 (A), human $G_{\alpha 13}$ transfected: sc-115367 (B) and sc-125358 (C) 293T whole cell lysates. Detection reagent used: m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM.



$G_{\alpha 13}$ (A-6): sc-518113. Western blot analysis of $G_{\alpha 13}$ expression in HEK293T whole cell lysate. Detection reagent used: m-IgG Fc BP-HRP: sc-525409.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.