

LF (BAL0106): sc-52055

BACKGROUND

Bacillus anthracis, a gram positive bacterium, is capsulogenic and toxinogenic and is the causative agent of anthrax. *Bacillus anthracis* secretes two toxins, which are composed of three proteins: the protective antigen (PA), the lethal factor (LF) and the edema factor (EF). PA and LF comprise a lethal toxin, which provokes a sudden death in animals, whereas the edema toxin, comprised of PA and EF, induces edema. The edema and the lethal factors are internalized into the target cells via the protective antigen. PA is the target-cell binding protein and is common to the two effector molecules, LF and EF, which exert their toxic effects once they are translocated to the cytosol by PA. PA is the major component of vaccines against anthrax since it confers protective immunity. The large-scale production of recombinant protein-based anthrax vaccines requires overexpression of the PA protein. LF plays an important role in the pathogenesis of anthrax. In addition, EF and LF exert adenylate cyclase and metalloprotease activity, respectively.

REFERENCES

1. Brossier, F., Guidi-Rontani, C. and Mock, M. 1998. Anthrax toxins. C.R. Seances Soc. Biol. Fil. 3: 437-444.
2. Brossier, F., Weber-Levy, M., Mock, M. and Sirard, J.C. 2000. Protective antigen-mediated antibody response against a heterologous protein produced *in vivo* by *Bacillus anthracis*. Infect. Immun. 10: 5731-5734.
3. Chauhan, V., Singh, A., Waheed, S.M., Singh, S. and Bhatnagar, R. 2001. Constitutive expression of protective antigen gene of *Bacillus anthracis* in *Escherichia coli*. Biochem. Biophys. Res. Commun. 2: 308-315.
4. Watters, J.W. and Dietrich, W.F. 2001. Genetic, physical, and transcript map of the *ltxs1* region of mouse chromosome 11. Genomics 2: 223-231.
5. Zhang, Y., Kida, Y., Kuwano, K., Misumi, Y., Ikehara, Y. and Arai, S. 2001. Role of furin in delivery of a CTL epitope of an anthrax toxin-fusion protein. Microbiol. Immunol. 2: 119-125.
6. Chang, H.H., Tsai, M.F., Chung, C.P., Chen, P.K., Hu, H.I., Kau, J.H., Huang, H.H., Lin, H.C. and Sun, D.S. 2006. Single-step purification of recombinant anthrax lethal factor from periplasm of *Escherichia coli*. J. Biotechnol. 126: 277-285.
7. Jiao, G.S., Simo, O., Nagata, M., O'Malley, S., Hemscheidt, T., Cregar, L., Millis, S.Z., Goldman, M.E. and Tang, C. 2006. Selectively guanidylated derivatives of neamine. Syntheses and inhibition of anthrax lethal factor protease. Bioorg. Med. Chem. Lett. 16: 5183-5189.
8. Sapa, R., Gaucher, S.P., Lachmann, J.S., Buffleben, G.M., Chirica, G.S., Comer, J.E., Peterson, J.W., Chopra, A.K. and Singh, A.K. 2006. Proteomic analyses of murine macrophages treated with *Bacillus anthracis* lethal toxin. Microb. Pathog. 41: 157-167
9. Mabry, R., Brasky, K., Geiger, R., Carrion, R., Jr., Hubbard, G.B., Leppla, S., Patterson, J.L., Georgiou, G. and Iverson, B.L. 2006. Detection of anthrax toxin in the serum of animals infected with *Bacillus anthracis* by using engineered immunoassays. Clin. Vaccine Immunol. 13: 671-677.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

LF (BAL0106) is a mouse monoclonal antibody raised against highly purified lethal factor (LF) antigen of *Bacillus anthracis*.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

LF (BAL0106) is recommended for detection of lethal factor antigen of *Bacillus anthracis* origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of LF: 101 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.