

PA (BAP0106): sc-52128

BACKGROUND

Bacillus anthracis, a Gram-positive bacterium, is capsulogenic and toxinogenic and is the causative agent of anthrax. *Bacillus anthracis* secretes two toxins, which are composed of three proteins: the protective antigen (PA), the lethal factor (LF) and the edema factor (EF). PA and LF comprise a lethal toxin, which provokes a sudden death in animals, whereas the edema toxin, comprised of PA and EF, induces edema. The edema and the lethal factors are internalized into the target cells via the protective antigen. PA is the target-cell binding protein and is common to the two effector molecules, LF and EF, which exert their toxic effects once they are translocated to the cytosol by PA. PA is the major component of vaccines against anthrax since it confers protective immunity. The large-scale production of recombinant protein-based anthrax vaccines requires overexpression of the PA protein. LF plays an important role in the pathogenesis of anthrax. In addition, EF and LF exert adenylate cyclase and metalloprotease activity, respectively.

REFERENCES

1. Brossier, F., Guidi-Rontani, C. and Mock, M. 1998. Anthrax toxins. C. R. Seances Soc. Biol. Fil. 3: 437-444.
2. Brossier, F., Weber-Levy, M., Mock, M. and Sirard, J.C. 2000. Protective antigen-mediated antibody response against a heterologous protein produced *in vivo* by *Bacillus anthracis*. Infect. Immun. 10: 5731-5734.
3. Chauhan, V., Singh, A., Waheed, S.M., Singh, S. and Bhatnagar, R. 2001. Constitutive expression of protective antigen gene of *Bacillus anthracis* in *Escherichia coli*. Biochem. Biophys. Res. Commun. 2: 308-315.
4. Zhang, Y., Kida, Y., Kuwano, K., Misumi, Y., Ikehara, Y. and Arai, S. 2001. Role of furin in delivery of a CTL epitope of an anthrax toxin-fusion protein. Microbiol. Immunol. 2: 119-125.
5. Watters, J.W. and Dietrich, W.F. 2001. Genetic, physical, and transcript map of the Itxs1 region of mouse chromosome 11. Genomics 2: 223-231.
6. Biagini, R.E., Sammons, D.L., Smith, J.P., MacKenzie, B.A., Striley, C.A., Snawder, J.E., Robertson, S.A. and Quinn, C.P. 2006. Rapid, sensitive, and specific lateral-flow immunochromatographic device to measure anti-anthrax protective antigen immunoglobulin γ in serum and whole blood. Clin. Vaccine Immunol. 13:541-546.
7. Fang, H., Xu, L., Chen, T.Y., Cyr, J.M. and Frucht, D.M. 2006. Anthrax lethal toxin has direct and potent inhibitory effects on B cell proliferation and immunoglobulin production. J. Immunol. 176: 6155-6161.
8. Sloat, B.R. and Cui, Z. 2006. Evaluation of the immune response induced by a nasal anthrax vaccine based on the protective antigen protein in anaesthetized and non-anaesthetized mice. J. Pharm. Pharmacol. 58: 439-447.
9. Rivera, J., Nakouzi, A., Abboud, N., Revskaya, E., Goldman, D., Collier, R.J., Dadachova, E. and Casadevall, A. 2006. A monoclonal antibody to *Bacillus anthracis* protective antigen defines a neutralizing epitope in domain 1. Infect. Immun. 74: 4149-4156.

SOURCE

PA (BAP0106) is a mouse monoclonal antibody raised against highly purified protective antigen (PA) of *Bacillus anthracis*.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PA (BAP0106) is recommended for detection of protective antigen of *Bacillus anthracis* origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of PA: 96 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.