

# IFN- $\gamma$ (XMG1.2): sc-52557

## BACKGROUND

Interferon (IFN)- $\gamma$  is an antiviral and antiparasitic agent produced by CD4<sup>+</sup>/CD8<sup>+</sup> lymphocytes and natural killer cells that undergo activation by antigens, mitogens or alloantigens. IFN- $\gamma$  production modulates T cell growth and differentiation and inhibits the growth of B cells. Synthesis of IFN- $\gamma$  is inducible by IL-2, FGF and EGF. The active form of IFN- $\gamma$  is a homodimer with each subunit containing six helices. The dimeric structure of human IFN- $\gamma$  is stabilized by non-covalent interactions through the interface of the helices. IFN- $\gamma$  translated precursor is 166 amino acids, including the 23 amino acid secretory sequence. Multiple forms exist due to variable glycosylation and under non-denaturing conditions due to dimers and tetramers.

## REFERENCES

1. Young, H.A. and Hardy, K.J. 1995. Role of interferon- $\gamma$  in immune cell regulation. *J. Leukoc. Biol.* 58: 373-381.
2. Dinarello, C.A., et al. 1998. Overview of interleukin-18: more than an interferon- $\gamma$  inducing factor. *J. Leukoc. Biol.* 63: 658-664.
3. Okamura, H., et al. 1998. Regulation of interferon- $\gamma$  production by IL-12 and IL-18. *Curr. Opin. Immunol.* 10: 259-264.
4. Costa-Pereira, A.P., et al. 2002. The antiviral response to  $\gamma$  interferon. *J. Virol.* 76: 9060-9068.
5. Zika, E., et al. 2003. Histone deacetylase 1/mSin3A disrupts  $\gamma$  interferon-induced CIITA function and major histocompatibility complex class II enhanceosome formation. *Mol. Cell. Biol.* 23: 3091-3102.
6. Schroder, K., et al. 2004. Interferon- $\gamma$ : an overview of signals, mechanisms and functions. *J. Leukoc. Biol.* 75: 163-189.
7. Ellis, T.N. and Beaman, B.L. 2004. Interferon- $\gamma$  activation of polymorphonuclear neutrophil function. *Immunology* 112: 2-12.
8. Sizemore, N., et al. 2004. Inhibitor of  $\kappa$ B kinase is required to activate a subset of interferon- $\gamma$  -stimulated genes. *Proc. Natl. Acad. Sci. USA* 101: 7994-7998.
9. Halfter, U.M., et al. 2005. Interferon- $\gamma$ -dependent tyrosine phosphorylation of MEKK4 via Pyk2 is regulated by Annexin II and SHP2 in keratinocytes. *Biochem. J.* 388: 17-28.

## CHROMOSOMAL LOCATION

Genetic locus: Ifng (mouse) mapping to 10 D2.

## SOURCE

IFN- $\gamma$  (XMG1.2) is a rat monoclonal antibody raised against recombinant IFN- $\gamma$  of mouse origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IFN- $\gamma$  (XMG1.2) is available conjugated fluorescein (sc-52557 FITC, 100 tests in 2 ml), for WB (RGB), IF, IHC(P) and FCM.


## APPLICATIONS

IFN- $\gamma$  (XMG1.2) is recommended for detection of IFN- $\gamma$  of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and flow cytometry (1  $\mu$ g per 1 x 10<sup>6</sup> cells).

Suitable for use as control antibody for IFN- $\gamma$  siRNA (m): sc-39607, IFN- $\gamma$  shRNA Plasmid (m): sc-39607-SH and IFN- $\gamma$  shRNA (m) Lentiviral Particles: sc-39607-V.

Molecular Weight of IFN- $\gamma$ : 20-25 kDa.

## DATA



IFN- $\gamma$  (XMG1.2): sc-52557. Western blot analysis of mouse recombinant IFN- $\gamma$ .

## SELECT PRODUCT CITATIONS

1. Copin, R., et al. 2007. MyD88-dependent activation of B220<sup>+</sup>CD11b<sup>+</sup>LY-6C<sup>+</sup> dendritic cells during *Brucella melitensis* infection. *J. Immunol.* 178: 5182-5191.
2. Aebischer, J., et al. 2011. IFN- $\gamma$  triggers a LIGHT-dependent selective death of motoneurons contributing to the non-cell-autonomous effects of mutant SOD1. *Cell Death Differ.* 18: 754-768.
3. Matsui, K., et al. 2015. Langerhans cell-like dendritic cells stimulated with an adjuvant direct the development of Th1 and Th2 cells *in vivo*. *Clin. Exp. Immunol.* 182: 101-107.
4. Wu, S., et al. 2016. Anti-inflammatory effects of *Boletus edulis* polysaccharide on asthma pathology. *Am. J. Transl. Res.* 8: 4478-4489.
5. Li, J., et al. 2017. VSIG4 inhibits proinflammatory macrophage activation by reprogramming mitochondrial pyruvate metabolism. *Nat. Commun.* 8: 1322.
6. Wu, S., et al. 2017. Anti-asthmatic effect of pitavastatin through aerosol inhalation is associated with CD4<sup>+</sup> CD25<sup>+</sup> Foxp3<sup>+</sup> T cells in an asthma mouse model. *Sci. Rep.* 7: 6084.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.