



JAM-A (M.Ab.F11): sc-52690

BACKGROUND

Junctional adhesion molecule (JAM) is a member of the immunoglobulin superfamily expressed in tight junctions of epithelial cells and endothelial cells. It is implicated in transendothelial migration of leukocytes. JAM is constitutively expressed on circulating monocytes, neutrophils, lymphocyte subsets and platelets. The JAM family consists of JAM-A, JAM-B and JAM-C, alternatively designated JAM-1, JAM-2 and JAM-3, respectively. JAM-A localizes with F-Actin at the cell-cell contacts and at the membrane ruffles. It is involved in cell to cell adhesion through homophilic interactions and plays a role in the organization of tight junctions and modulation of leukocyte extravasation. JAM-B interacts with discrete subsets of PBLs, suggesting that it may play a role in lymphocyte trafficking. JAM-B and JAM-C proteins are binding partners; JAM-C may be a functional JAM-B receptor. Specifically, JAM-B adheres to T cells through heterotypic interactions with JAM-C. The JAM-B/JAM-C interaction may play a role in T, NK and dendritic cellular inflammation.

REFERENCES

1. Martin-Padura, I., et al. 1998. Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J. Cell Biol.* 142: 117-127.
2. Ozaki, H., et al. 1999. Cutting edge: combined treatment of TNF α and IFN- γ causes redistribution of junctional adhesion molecule in human endothelial cells. *J. Immunol.* 163: 553-557.
3. Ozaki, H., et al. 2000. Junctional adhesion molecule (JAM) is phosphorylated by protein kinase C upon platelet activation. *Biochem. Biophys. Res. Commun.* 276: 873-878.
4. Ebnet, K., et al. 2000. Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6 and ZO-1. *J. Biol. Chem.* 275: 27979-27988.
5. Dejana, E., et al. 2000. The molecular organization of endothelial junctions and their functional role in vascular morphogenesis and permeability. *Int. J. Dev. Biol.* 44: 743-748.
6. Bazzoni, G., et al. 2000. Homophilic interaction of junctional adhesion molecule. *J. Biol. Chem.* 275: 30970-30976.
7. Arrate, M.P., et al. 2001. Cloning of human junctional adhesion molecule 3 (JAM-3) and its identification as the JAM-2 counter-receptor. *J. Biol. Chem.* 276: 45826-45832.
8. Liang, T.W., et al. 2002. Vascular endothelial-junctional adhesion molecule (VE-JAM)/JAM-2 interacts with T, NK, and dendritic cells through JAM-3. *J. Immunol.* 168: 1618-1626.

CHROMOSOMAL LOCATION

Genetic locus: F11R (human) mapping to 1q23.3.

SOURCE

JAM-A (M.Ab.F11) is a mouse monoclonal antibody raised against JAM-A of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


JAM-A (M.Ab.F11) is recommended for detection of JAM-A of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and flow cytometry (1 μ g per 1×10^6 cells).

Suitable for use as control antibody for JAM-A siRNA (h): sc-43139, JAM-A shRNA Plasmid (h): sc-43139-SH and JAM-A shRNA (h) Lentiviral Particles: sc-43139-V.

Molecular Weight of JAM-A: 36 kDa.

Positive Controls: human platelet extract: sc-363773 or T84 whole cell lysate: sc-364797.

DATA



JAM-A (M.Ab.F11): sc-52690. Indirect FCM analysis of human peripheral blood leukocytes stained with JAM-A (M.Ab.F11), followed by PE-conjugated goat anti-mouse IgG: sc-3738. Black line histogram represents the isotype control, normal mouse IgG₁: sc-3877.

SELECT PRODUCT CITATIONS

1. Terral, G., et al. 2017. Epitope characterization of anti-JAM-A antibodies using orthogonal mass spectrometry and surface plasmon resonance approaches. *MAbs* 9: 1317-1326.
2. Zhang, H. and Lu, W. 2018. LncRNA SNHG12 regulates gastric cancer progression by acting as a molecular sponge of miR-320. *Mol. Med. Rep.* 17: 2743-2749.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

CONJUGATES

See **JAM-A (J10.4): sc-53623** for JAM-A antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.