

TEM8 (200C1339): sc-52959

BACKGROUND

The tripartite toxin secreted by *Bacillus anthracis* is the causative agent of anthrax evading the immune system and killing the host during a systemic infection. Two components of the toxin, edema factor (EF) and lethal factor (LF), enzymatically modify substrates within the cytosol of mammalian cells. The third component, protective antigen (PA), binds to a cellular receptor, designated ATR (anthrax toxin receptor), which mediates the delivery of the enzymatic components to the cytosol. TEM8 (tumor endothelial marker 8) is one of the tumor-specific endothelial markers (TEMs) whose N-terminus encodes ATR. TEM8 is highly expressed in tumor endothelial cells but not in normal endothelial cells. TEMs have elevated expression during tumor angiogenesis. Four TEM genes, TEM1, TEM5, TEM7 and TEM8, encode the TEM proteins, which contain putative transmembrane domains. ATR is a type I membrane protein with an extracellular von Willebrand factor A domain that binds directly to PA. The first 364 amino acids of ATR protein are identical to those of TEM8. However, the C-terminal ends of the ATR and TEM8 proteins are different, presumably due to alternative splicing. A soluble version of von Willebrand factor A domain seems to protect cells from the toxin action.

REFERENCES

1. Leppla, S.H. 1982. Anthrax toxin edema factor: a bacterial adenylate cyclase that increases cAMP concentration in eukaryotic cells. *Proc. Natl. Acad. Sci. USA* 79: 3162-3166.
2. O'Brien, J., et al. 1985. Effects of anthrax toxin components on human neutrophils. *Infect. Immun.* 47: 306-310.
3. Duesbery, N.S., et al. 1998. Proteolytic inactivation of MAP-kinase-kinase by anthrax lethal factor. *Science* 280: 734-737.
4. Pellizzari, R., et al. 1999. Anthrax lethal factor cleaves MKK3 in macrophages and inhibits the LPS/IFN γ -induced release of NO and TNF α . *FEBS Lett.* 462: 199-204.
5. St Croix, B., et al. 2000. Genes expressed in human tumor endothelium. *Science* 289: 1197-1202.
6. Bradley, K.A., et al. 2001. Identification of the cellular receptor for anthrax toxin. *Nature* 414: 225-229.
7. Carson-Walter, E.B., et al. 2001. Cell surface tumor endothelial markers are conserved in mice and humans. *Cancer Res.* 61: 6649-6655.

CHROMOSOMAL LOCATION

Genetic locus: ANTXR1 (human) mapping to 2p13.3.

SOURCE

TEM8 (200C1339) is a mouse monoclonal antibody raised against amino acids 481-497 of TEM8 of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


TEM8 (200C1339) is recommended for detection of TEM8 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for TEM8 siRNA (h): sc-44144, TEM8 shRNA Plasmid (h): sc-44144-SH and TEM8 shRNA (h) Lentiviral Particles: sc-44144-V.


Molecular Weight of TEM8: 63 kDa.

Positive Controls: Raji whole cell lysate: sc-364236, HeLa whole cell lysate: sc-2200 or human PBL whole cell lysate.

DATA



TEM8 (200C1339): sc-52959. Western blot analysis of TEM8 expression in HeLa (A) and Raji (B) whole cell lysates.



TEM8 (200C1339): sc-52959. Western blot analysis of TEM8 expression in human PBL whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.