

# Ig $\kappa$ chain (OX20): sc-53080

## BACKGROUND

Antibody producing cells of the immune system require multiple rearrangements of immunoglobulin (antibody, Ig) genes. Immunoglobulins are four-chain, Y-shaped, monomeric structures of two identical heavy chains and two identical light chains held together through interchain disulfide bonds. Immunoglobulins in vertebrates help to remove non-self molecules or cells (antigens) by recognizing and binding to the antigen and carrying out effector functions that activate the immune system. Variable genetic combinations of the five heavy chain classes (M, D, G, E and A) and the two light chain isotypes,  $\kappa$  and  $\lambda$ , confer the role of an antibody. The variable region genes encoding immunoglobulin  $\kappa$  and  $\lambda$  chains are assembled from three DNA segments, the V, C and J genes. Human  $\kappa$  light chain genes map to chromosome 2 and the human  $\lambda$  light chain genes map to chromosome 22.  $\kappa$  gene recombination can precede  $\lambda$  gene recombination during B cell ontogeny and only a single light chain type is expressed in individual B cells. Antibodies in camels and sharks can lack light chains, suggesting that light chains may not be essential for antigen binding in some vertebrates.

## REFERENCES

- Hieter, P.A., et al. 1980. Cloned human and mouse  $\kappa$  immunoglobulin constant and J region genes conserve homology in functional segments. *Cell* 22: 197-207.
- Mason, D.W., et al. 1981. The rat mixed lymphocyte reaction: roles of a dendritic cell in intestinal lymph and T cell subsets defined by monoclonal antibodies. *Immunology* 44: 75-87.
- Dyer, M.J., et al. 1981. Committed T lymphocyte stem cells of rats. Characterization by surface W3/13 antigen and radiosensitivity. *J. Exp. Med.* 154: 1164-1177.
- Hieter, P.A., et al. 1982. Evolution of human immunoglobulin  $\kappa$  J region genes. *J. Biol. Chem.* 257: 1516-1522.
- Durdik, J., et al. 1984. Novel  $\kappa$  light chain gene rearrangements in mouse  $\lambda$  light chain-producing B lymphocytes. *Nature* 307: 749-752.
- Horejsi, V., et al. 1986. Monoclonal antibodies against human leukocyte antigens. I. Antibodies against  $\beta_2$ -Microglobulin, immunoglobulin  $\kappa$  light chains, HLA-DR-like antigens, T8 antigen, T1 antigen, a monocyte antigen, and a pan-leukocyte antigen. *Folia Biol.* 32: 12-25.
- Pilstrom, L. 2002. The mysterious immunoglobulin light chain. *Dev. Comp. Immunol.* 26: 207-215.
- Li, M., et al. 2004. Expression of immunoglobulin  $\kappa$  light chain constant region in abnormal human cervical epithelial cells. *Int. J. Biochem. Cell Biol.* 36: 2250-2257.
- LocusLink Report (LocusID: 3514). <http://www.ncbi.nlm.nih.gov/LocusLink/>

## CHROMOSOMAL LOCATION

Genetic locus: Igkc (mouse) mapping to 6 C1.

## SOURCE

Ig  $\kappa$  chain (OX20) is a rat monoclonal antibody raised against purified IgG<sub>2</sub> of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>2</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Ig  $\kappa$  chain (OX20) is available conjugated to agarose (sc-53080 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-53080 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-53080 PE), fluorescein (sc-53080 FITC), Alexa Fluor<sup>®</sup> 488 (sc-53080 AF488), Alexa Fluor<sup>®</sup> 546 (sc-53080 AF546), Alexa Fluor<sup>®</sup> 594 (sc-53080 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-53080 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-53080 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-53080 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA


## APPLICATIONS

Ig  $\kappa$  chain (OX20) is recommended for detection of Ig  $\kappa$  light chains of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and flow cytometry (1  $\mu$ g per 1 x 10<sup>6</sup> cells).


Molecular Weight of Ig  $\kappa$  chain: 28 kDa.

Positive Controls: mouse PBL whole cell lysate.

## DATA



Ig  $\kappa$  chain (OX20): sc-53080. Western blot analysis of Ig  $\kappa$  chain expression in mouse PBL whole cell lysate.



Ig  $\kappa$  chain (OX20) HRP: sc-53080 HRP. Direct western blot analysis of Ig  $\kappa$  chain expression in mouse PBL whole cell lysate.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.