

# cyclin A (E70.1): sc-53231

## BACKGROUND

The critical role that the family of regulatory proteins known as cyclins play in eukaryotic cell cycle regulation is well established. The best-characterized cyclin complex is the mitotic cyclin B/Cdc2 p34 kinase, the active component of maturing promoting factor. Cyclin A accumulates prior to cyclin B in the cell cycle, appears to be involved in control of S phase and has been shown to associate with cyclin-dependent kinase-2 (Cdk2). In addition, cyclin A has been implicated in cell transformation and is found in complexes with E1A, transcription factors DRTF1 and E2F and retinoblastoma protein, p110. A second form of cyclin A, named cyclin A1 because of its high sequence homology to *Xenopus* cyclin A1, is most highly expressed in germ cells. It has been proposed that cyclin A1 can associate with Cdk2, p39 and Cdc2 p34.

## REFERENCES

1. Draetta, G., et al. 1989. Cdc2 protein kinase is complexed with both cyclin A and B: evidence for proteolytic inactivation of MPF. *Cell* 56: 829-838.
2. Giordano, A., et al. 1989. A 60 kDa Cdc2-associated polypeptide complexes with the E1A proteins in adenovirus-infected cells. *Cell* 58: 981-990.
3. Gautier, J., et al. 1990. Cyclin is a component of maturation-promoting factor from *Xenopus*. *Cell* 60: 487-494.
4. Wang, J., et al. 1990. Hepatitis B virus integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 343: 555-557.
5. Pines, J., et al. 1990. Human cyclin A is adenovirus E1A-associated protein p60 and behaves differently from cyclin B. *Nature* 346: 760-763.
6. Bandara, L.R., et al. 1991. Cyclin A and the retinoblastoma gene product complex with a common transcription factor. *Nature* 352: 249-251.
7. Williams, R.T., et al. 1992. Co-purification of p34Cdc2/p58 cyclin A proline-directed protein kinase and the retinoblastoma tumor susceptibility gene product: interaction of an oncogenic Serine/Threonine protein kinase with a tumor-suppressor protein. *Oncogene* 7: 423-432.
8. Lees, E., et al. 1992. Cyclin E/Cdk2 and cyclin A/Cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev.* 6: 1874-1885.
9. Yang, R., et al. 1997. Characterization of a second human cyclin A that is highly expressed in testis and in several leukemic cell lines. *Cancer Res.* 57: 913-930.

## CHROMOSOMAL LOCATION

Genetic locus: CCNA2 (human) mapping to 4q27; Ccna2 (mouse) mapping to 3 B.

## SOURCE

cyclin A (E70.1) is a mouse monoclonal antibody raised against cyclin A of bovine origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

cyclin A (E70.1) is recommended for detection of cyclin A of mouse, rat, human and bovine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for cyclin A siRNA (h): sc-29282, cyclin A siRNA (m): sc-29283, cyclin A shRNA Plasmid (h): sc-29282-SH, cyclin A shRNA Plasmid (m): sc-29283-SH, cyclin A shRNA (h) Lentiviral Particles: sc-29282-V and cyclin A shRNA (m) Lentiviral Particles: sc-29283-V.


Molecular Weight of cyclin A: 54 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, HuT 78 whole cell lysate: sc-2208 or F9 cell lysate: sc-2245.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



cyclin A (E70.1): sc-53231. Western blot analysis of cyclin A expression in K-562 (A) and F9 (B) whole cell lysates.



cyclin A (E70.1): sc-53231. Western blot analysis of cyclin A expression in K-562 (A) and HuT 78 (B) whole cell lysates.

## SELECT PRODUCT CITATIONS

1. Song, X., et al. 2018. SRC-3 inhibition blocks tumor growth of pancreatic ductal adenocarcinoma. *Cancer Lett.* 442: 310-319.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **cyclin A (B-8): sc-271682** for cyclin A antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.