

Cytokeratin 15/18 (LC18N): sc-53254

BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins are useful markers of tissue differentiation and, therefore, the characterization of malignant tumors. Cytokeratin 15 is an intermediate filament (IF) type I protein that is responsible for the mechanical integrity of epithelial cells. Cytokeratin 15 is a specific marker of stem cells of the hair-follicle bulge and may be a useful marker for diagnosis between basal cell carcinoma and trichoepithelioma. Cytokeratin 18 is involved in keratin organization and distribution. Phosphorylation of Ser 18 of Cytokeratin 18 is essential for its association with 14-3-3 proteins. Cytokeratin 18 is expressed in a majority of adenocarcinomas and basal cell carcinomas.

REFERENCES

- Linder, S., Havelka, A.M., Ueno, T. and Shoshan, M.C. 2004. Determining tumor apoptosis and necrosis in patient serum using Cytokeratin 18 as a biomarker. *Cancer Lett.* 214: 1-9.
- Roth, G.A., Krenn, C., Brunner, M., Moser, B., Ploder, M., Spittler, A., Pelinka, L., Sautner, T., Wolner, E., Boltz-Nitulescu, G. and Ankersmit, H.J. 2004. Elevated serum levels of epithelial cell apoptosis-specific Cytokeratin 18 neopeptide m30 in critically ill patients. *Shock* 22: 218-220.
- Viswanathan, V.K., Lukic, S., Koutsouris, A., Miao, R., Muza, M.M. and Hecht, G. 2004. Cytokeratin 18 interacts with the enteropathogenic *Escherichia coli* secreted protein F (EspF) and is redistributed after infection. *Cell. Microbiol.* 6: 987-997.
- Kronenberger, B., Wagner, M., Herrmann, E., Mihm, U., Piiper, A., Sarrazin, C. and Zeuzem, S. 2005. Apoptotic Cytokeratin 18 neopeptides in serum of patients with chronic hepatitis C. *J. Viral Hepat.* 12: 307-314.
- Sharma, A.D., Cantz, T., Richter, R., Eckert, K., Henschler, R., Wilkens, L., Jochheim-Richter, A., Arseniev, L. and Ott, M. 2005. Human cord blood stem cells generate human Cytokeratin 18-negative hepatocyte-like cells in injured mouse liver. *Am. J. Pathol.* 167: 555-564.
- Yaniw, D. and Hu, J. 2005. Epithelium-specific ets transcription factor 2 upregulates Cytokeratin 18 expression in pulmonary epithelial cells through an interaction with Cytokeratin 18 intron 1. *Cell Res.* 15: 423-429.
- Cianga, P., Cianga, C., Anisie, E., Zlei, M., Ungureanu, D., Munteanu, D., Chifu, C., Diaconu, C. and Carasevici, E. 2006. High values of soluble Cytokeratin 18 in breast carcinomas may have different meanings. *Rev. Med. Chir. Soc. Med. Nat. Iasi* 109: 836-840.
- Pichon, M.F., Labroquere, M., Rezai, K. and Lokiec, F. 2006. Variations of soluble fas and Cytokeratin 18-Asp 396 neo-epitope in different cancers during chemotherapy. *Anticancer Res.* 26: 2387-2392.
- Yim, H., Kim, J.E., Shin, J.Y., Ye, Y.M., Park, H.S. and Nahm, D.H. 2006. Antigen-binding characteristics of circulating IgG autoantibodies to Cytokeratin 18 protein in patients with nonallergic asthma. *J. Korean Med. Sci.* 21: 652-655.

CHROMOSOMAL LOCATION

Genetic locus: KRT15 (human) mapping to 17q21.2, KRT18 (human) mapping to 12q13.13.

SOURCE

Cytokeratin 15/18 (LC18N) is a mouse monoclonal antibody raised against an N-terminal synthetic peptide of Cytokeratin 18 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Cytokeratin 15/18 (LC18N) is recommended for detection of Cytokeratin 15/18 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of Cytokeratin 15: 52 kDa.

Molecular Weight of Cytokeratin 18: 45 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201.

SELECT PRODUCT CITATIONS

- Ma, X., Chen, S., Jin, W. and Gao, Y. 2017. Th1/Th2 PB balance and CD200 expression of patients with active severe alopecia areata. *Exp. Ther. Med.* 13: 2883-2887.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Cytokeratin 18 (RGE53): sc-32329** for Cytokeratin 18 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.