

# ErbB-4 (HFR1): sc-53280

## BACKGROUND

The EGF receptor family comprises several related receptor tyrosine kinases that are frequently overexpressed in a variety of carcinomas. Members of this receptor family include EGFR (HER1), Neu (ErbB-2, HER2), ErbB-3 (HER3) and ErbB-4 (HER4), which form either homodimers or heterodimers upon ligand binding. The gene encoding ErbB-4 is expressed as a full length protein, which produces a short membrane-anchored cytoplasmic domain fragment and a long ectodomain fragment. The short fragment is heavily tyrosine phosphorylated and possesses tyrosine kinase catalytic activity toward an exogenous substrate. Proteolytic cleavage of ErbB-4 is promoted by the binding of heregulin. ErbB-4 is involved in cell proliferation and differentiation and its expression is highest in breast carcinoma cell lines, normal skeletal muscle, heart, pituitary, brain and cerebellum.

## REFERENCES

1. Plowman, G.D., et al. 1993. Ligand-specific activation of HER4/p180ErbB-4, a fourth member of the epidermal growth factor receptor family. *Proc. Natl. Acad. Sci. USA* 90: 1746-1750.
2. Zimonjic, D.B., et al. 1995. Localization of the human HER4/ErbB-4 gene to chromosome 2. *Oncogene* 10: 1235-1237.

## CHROMOSOMAL LOCATION

Genetic locus: ERBB4 (human) mapping to 2q34; ErbB4 (mouse) mapping to 1 C3.

## SOURCE

ErbB-4 (HFR1) is a mouse monoclonal antibody raised against amino acids 1249-1264 of ErbB-4 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

ErbB-4 (HFR1) is recommended for detection of ErbB-4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for ErbB-4 siRNA (h): sc-35329, ErbB-4 siRNA (m): sc-35330, ErbB-4 shRNA Plasmid (h): sc-35329-SH, ErbB-4 shRNA Plasmid (m): sc-35330-SH, ErbB-4 shRNA (h) Lentiviral Particles: sc-35329-V and ErbB-4 shRNA (m) Lentiviral Particles: sc-35330-V.

Molecular Weight of ErbB-4 precursor: 180 kDa.


Molecular Weight of ErbB-4 cleaved forms: 80/120 kDa.

Positive Controls: mouse brain extract: sc-2253 or ErbB-4 transfected NIH/3T3 whole cell lysate.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



ErbB-4 (HFR1): sc-53280. Western blot analysis of ErbB-4 expression in ErbB-4 transfected NIH/3T3 whole cell lysate.



ErbB-4 (HFR1): sc-53280. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic and nuclear staining of myocytes.

## SELECT PRODUCT CITATIONS

1. Eckert, J.M., et al. 2009. Neuregulin-1 β and neuregulin-1 α differentially affect the migration and invasion of malignant peripheral nerve sheath tumor cells. *Glia* 57: 1501-1520.
2. Levchenko, V., et al. 2010. EGF and its related growth factors mediate sodium transport in mpkCCDC14 cells via ErbB2 (neu/HER-2) receptor. *J. Cell. Physiol.* 223: 252-259.
3. Biltekin, B., et al. 2023. *In vitro* effects of heparin-binding epidermal growth factor on adhesion stage of implantation. *Rom. J. Morphol. Embryol.* 64: 493-500.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **ErbB-4 (C-7): sc-8050** for ErbB-4 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.