

# m-IgG Fc BP-CFL 790: sc-533658

## BACKGROUND

Mouse IgG Fc binding protein (m-IgG Fc BP) conjugated to CruzFluor™ 790 (CFL 790) is a strongly recommended alternative to conventional goat/rabbit anti-mouse IgG secondary antibodies for NIR Western Blotting (WB), immunofluorescence (IF) and flow cytometry (FCM) signal enhancement. CruzFluor™ 790 (CFL 790) is an infrared fluorescent dye that is an excellent substitute for AlexaFluor® 790, offering comparable photostability and the ability to resist protein quenching. Suitable for use with Near-Infrared (NIR) imaging systems, such as LI-COR/Odyssey, Invitrogen/iBright and other comparable systems. Mouse IgG Fc binding protein is a highly specific reagent that provides strong signal with minimal background and virtually complete elimination of lot to lot variation associated with conventionally generated secondary antibodies. Mouse IgG Fc binding protein (m-IgG Fc BP) is suitable for binding to the Fc region of most, but not all, mouse IgG<sub>1</sub>, IgG<sub>2a</sub> and IgG<sub>2b</sub> immunoglobulins, and to a lesser extent to mouse IgG<sub>3</sub>; not suitable for use with mouse monoclonal IgM, IgA and IgE. Not cross reactive with human, rat, rabbit and goat IgG antibodies.

## SOURCE

m-IgG Fc BP-CFL 790 is a purified recombinant mouse IgG Fc light chain binding protein conjugated to CruzFluor™ 790 (CFL 790).

## PRODUCT

Each vial contains 50 µg mouse IgG Fc binding protein-CFL 790 in 0.5 ml of PBS containing 0.1% gelatin and 0.1% sodium azide.

## APPLICATIONS

m-IgG Fc BP-CFL 790 is recommended for detection of mouse IgG Fc by NIR Western Blotting (starting dilution: 1:1000, dilution range: 1:1000-1:10000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:200) and flow cytometry (0.5-1 µg per 1 x 10<sup>6</sup> cells). Optimal dilution to be determined by titration.

For Western Blotting using tissue extracts and m-IgG Fc BP-CFL 790, we strongly recommend subtracting endogenous immunoglobulins from extracts with Protein G PLUS-Agarose Reagent: sc-2002, to prevent Western Blotting interference when detecting proteins of approximately 25 kDa in size.

## RECOMMENDED SUPPORT PRODUCTS


- RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- TBS Blotting A, lyophilized powder in single-use bottle: sc-2333
- UltraCruz® PVDF Transfer Membrane, 0.45 µm, 30 cm x 3 m roll: sc-3723
- UltraCruz® Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- UltraCruz® Gel Incubation Trays, 100 per pack: sc-201755 (blue), sc-201756 (green), sc-201757 (pink), sc-201758 (yellow), sc-201759 (orange)

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



β-Actin (C4): sc-47778. Near-infrared western blot analysis of β-Actin expression in HeLa (A), MCF7 (B) and A-431 (C) whole cell lysates. Blocked with UltraCruz®. Blocking Reagent: sc-516214. Detection reagent used: m-IgG Fc BP-CFL 790: sc-533658.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CRUZFLUOR™ SPECTRAL PROPERTIES

PRODUCT	CAT. #	EXCITATION MAXIMUM	EMISSION MAXIMUM
m-IgG Fc BP-CFL 488	sc-533653	488 nm	514 nm
m-IgG <sub>1</sub> BP-CFL 488	sc-533661		
m-IgG Fc BP-CFL 555	sc-533654	556 nm	569 nm
m-IgG <sub>1</sub> BP-CFL 555	sc-533662		
m-IgG Fc BP-CFL 594	sc-533655	587 nm	603 nm
m-IgG <sub>1</sub> BP-CFL 594	sc-533663		
m-IgG Fc BP-CFL 647	sc-533656	654 nm	669 nm
m-IgG <sub>1</sub> BP-CFL 647	sc-533664		
m-IgG Fc BP-CFL 680	sc-533657	683 nm	700 nm
m-IgG <sub>1</sub> BP-CFL 680	sc-533665		
m-IgG Fc BP-CFL 790	sc-533658	786 nm	811 nm
m-IgG <sub>1</sub> BP-CFL 790	sc-533666		