

NGFR p75 (H-6): sc-55467

BACKGROUND

The Trk oncogene encodes a membrane-spanning protein tyrosine kinase, gp140Trk, whose expression is restricted *in vivo* to neurons of the sensory spinal and cranial ganglia of neural crest origin. Nerve growth factor (NGF) stimulates tyrosine phosphorylation of gp140Trk in neural cell lines and in embryonic dorsal root ganglia. Tyrosine phosphorylation of Trk by NGF is rapid, specific and occurs with picomolar quantities of factor, indicating that the response is mediated by physiological amounts of NGF, suggesting that gp140Trk participates in the primary signal transduction mechanism of NGF. An additional component of the gp140 Trk receptor complex, NGFR p75, binds to the neurotrophic factors with low affinity but is required for efficient signaling. NGFR p75 accelerates gp140Trk activation and may recruit downstream effector molecules to the liganded complex.

REFERENCES

1. Martin-Zanca, D., et al. 1986. A human oncogene formed by the fusion of truncated tropomyosin and protein tyrosine kinase sequences. *Nature* 319: 743-748.
2. Reinach, F.C., et al. 1986. Tissue-specific expression of the human tropomyosin gene involved in the generation of the Trk oncogene. *Nature* 322: 648-650.

CHROMOSOMAL LOCATION

Genetic locus: NGFR (human) mapping to 17q21.33; Ngfr (mouse) mapping to 11 D.

SOURCE

NGFR p75 (H-6) is a mouse monoclonal antibody raised against amino acids 29-165 of NGFR p75 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

NGFR p75 (H-6) is recommended for detection of NGFR p75 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NGFR p75 siRNA (h): sc-36058, NGFR p75 siRNA (m): sc-37268, NGFR p75 shRNA Plasmid (h): sc-36058-SH, NGFR p75 shRNA Plasmid (m): sc-37268-SH, NGFR p75 shRNA (h) Lentiviral Particles: sc-36058-V and NGFR p75 shRNA (m) Lentiviral Particles: sc-37268-V.


Molecular Weight of NGFR p75: 75 kDa.

Positive Controls: PC-12 cell lysate: sc-2250, SK-N-MC cell lysate: sc-2237 or mouse brain extract: sc-2253.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NGFR p75 (H-6): sc-55467. Western blot analysis of NGFR p75 expression in SK-N-MC whole cell lysate (A) and mouse brain tissue extract (B).

SELECT PRODUCT CITATIONS

1. Chavez-Valdez, R., et al. 2012. Effect of hyperoxic exposure during early development on neurotrophin expression in the carotid body and nucleus tractus solitarii. *J. Appl. Physiol.* 112: 1762-1772.
2. Di Cara, G., et al. 2013. Proteomic profiling of Trastuzumab (Herceptin®)-sensitive and -resistant SKBR-3 breast cancer cells. *Anticancer Res.* 33: 489-503.
3. Stucky, A., et al. 2016. Prenatal cocaine exposure upregulates BDNF-TrkB signaling. *PLoS ONE* 11: e0160585.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **NGFR p75 (B-1): sc-271708** for NGFR p75 antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.