

Cdk4 (DCS-31): sc-56277



The Power to Question

BACKGROUND

Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Several Cdk proteins have been identified, including Cdk2-Cdk8, PCTAIRE-1-PCTAIRE-3, PITALRE and PITSLRE. Cdk4, in complex with D-type cyclins, is thought to regulate cell growth during the G₁ phase of the cell cycle. This association with a D-type cyclin upregulates Cdk4 activity, whereas binding to the Cdk inhibitor p16 downregulates Cdk4 activity. Activation of the Cdk4-cyclin complexes requires phosphorylation on a single threonyl residue of Cdk4, catalyzed by a Cdk-activating protein (CAK).

CHROMOSOMAL LOCATION

Genetic locus: CDK4 (human) mapping to 12q14.1; Cdk4 (mouse) mapping to 10 D3.

SOURCE

Cdk4 (DCS-31) is a mouse monoclonal antibody raised against full length Cdk4 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Cdk4 (DCS-31) is available conjugated to agarose (sc-56277 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-56277 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-56277 PE), fluorescein (sc-56277 FITC), Alexa Fluor® 488 (sc-56277 AF488), Alexa Fluor® 546 (sc-56277 AF546), Alexa Fluor® 594 (sc-56277 AF594) or Alexa Fluor® 647 (sc-56277 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-56277 AF680) or Alexa Fluor® 790 (sc-56277 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Cdk4 (DCS-31) is recommended for detection of Cdk4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Cdk types.

Suitable for use as control antibody for Cdk4 siRNA (h): sc-29261, Cdk4 siRNA (m): sc-29262, Cdk4 shRNA Plasmid (h): sc-29261-SH, Cdk4 shRNA Plasmid (m): sc-29262-SH, Cdk4 shRNA (h) Lentiviral Particles: sc-29261-V and Cdk4 shRNA (m) Lentiviral Particles: sc-29262-V.


Molecular Weight of Cdk4: 34 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, MCF7 whole cell lysate: sc-2206 or MDA-MB-231 cell lysate: sc-2232.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Cdk4 (DCS-31): sc-56277. Western blot analysis of Cdk4 expression in HeLa (A), MCF7 (B), MDA-MB-231 (C) and ZR-75-1 (D) whole cell lysates. Detection reagent used: m-IgGκ BP-HRP: sc-516102.



Cdk4 (DCS-31): sc-56277. Western blot analysis of Cdk4 expression in untreated (A) and L-Mimosine (sc-201536) treated (B) MDA-MB-231 whole cell lysates. Note down regulation of Cdk4 expression in lane B.

SELECT PRODUCT CITATIONS

- Roger, P.P., et al. 1999. Nature of the critical labile event that controls RB phosphorylation in the cyclic AMP-dependent cell cycle of thyrocytes in primary culture. *Exp. Cell Res.* 252: 492-498.
- Depoortere, F., et al. 2000. Transforming growth factor β_1 selectively inhibits the cyclic AMP-dependent proliferation of primary thyroid epithelial cells by preventing the association of cyclin D3—Cdk4 with nuclear p27^{kip1}. *Mol. Biol. Cell* 11: 1061-1076.
- Wu, W., et al. 2009. Antibody array analysis with label-based detection and resolution of protein size. *Mol. Cell. Proteomics* 8: 245-257.
- Yan, J., et al. 2011. Andrographolide induces cell cycle arrest and apoptosis in human rheumatoid arthritis fibroblast-like synoviocytes. *Cell Biol. Toxicol.* 28: 47-56.
- Klaic, L., et al. 2012. Celastrol analogues as inducers of the heat shock response. Design and synthesis of affinity probes for the identification of protein targets. *ACS Chem. Biol.* 7: 928-937.
- Fischer, B.M., et al. 2013. Increased expression of senescence markers in cystic fibrosis airways. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 304: L394-L400.
- Han, Y.S., et al. 2015. Fucoidan inhibits the migration and proliferation of HT-29 human colon cancer cells via the phosphoinositide-3 kinase/Akt/mechanistic target of rapamycin pathways. *Mol. Med. Rep.* 12: 3446-3452.
- Colleoni, B., et al. 2017. JNKs function as Cdk4-activating kinases by phosphorylating Cdk4 and p21. *Oncogene* 36: 4349-4361.
- Liao, P., et al. 2017. Mutant p53 gains its function via c-Myc activation upon Cdk4 phosphorylation at serine 249 and consequent PIN1 binding. *Mol. Cell* 68: 1134-1146.

RESEARCH USE

For research use only, not for use in diagnostic procedures.