

N-Myc (NCM II 100): sc-56729

BACKGROUND

The v-Myc oncogene, initially identified in the MC29 avian retrovirus, causes myelocytomas, carcinomas, sarcomas and lymphomas, and belongs to a family of oncogenes conserved throughout evolution. In humans, the family consists of five genes: c-Myc, N-Myc, R-Myc, L-Myc and B-Myc. Amplification of the N-Myc gene has been found in human neuroblastomas and cell lines (3,5). The extent of N-Myc amplification correlates well with the stage of neuroblastoma disease. Immunological studies have shown that the human N-Myc gene encodes a nuclear phosphoprotein that exhibits relatively short (30 min) half life *in vivo*. The prototype member of the family, c-Myc p67, binds DNA in a sequence-specific manner subsequent to dimerization with a second basic region helix-loop-helix leucine zipper motif protein designated Max.

CHROMOSOMAL LOCATION

Genetic locus: MYCN (human) mapping to 2p24.3; Mycn (mouse) mapping to 12 A1.1.

SOURCE

N-Myc (NCM II 100) is a mouse monoclonal antibody raised against recombinant N-Myc of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

N-Myc (NCM II 100) is available conjugated to agarose (sc-56729 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-56729 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-56729 PE), fluorescein (sc-56729 FITC), Alexa Fluor® 488 (sc-56729 AF488), Alexa Fluor® 546 (sc-56729 AF546), Alexa Fluor® 594 (sc-56729 AF594) or Alexa Fluor® 647 (sc-56729 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-56729 AF680) or Alexa Fluor® 790 (sc-56729 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

N-Myc (NCM II 100) is recommended for detection of N-Myc encoded proteins and their cleavage products of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for N-Myc siRNA (h): sc-36003, N-Myc siRNA (m): sc-38087, N-Myc shRNA Plasmid (h): sc-36003-SH, N-Myc shRNA Plasmid (m): sc-38087-SH, N-Myc shRNA (h) Lentiviral Particles: sc-36003-V and N-Myc shRNA (m) Lentiviral Particles: sc-38087-V.

Molecular Weight of N-Myc: 67 kDa.

Positive Controls: N-Myc (m): 293T Lysate: sc-121906, IMR-32 cell lysate: sc-2409 or SH-SY5Y cell lysate: sc-3812.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



N-Myc (NCM II 100): sc-56729. Western blot analysis of N-Myc expression in IMR-32 (A), SH-SY5Y (B) and BE (2)-M17 (C) whole cell lysates.



N-Myc (NCM II 100): sc-56729. Western blot analysis of N-Myc expression in non-transfected: sc-117752 (A) and mouse N-Myc transfected: sc-121906 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Chen, H.F., et al. 2011. Surface marker epithelial cell adhesion molecule and E-cadherin facilitate the identification and selection of induced pluripotent stem cells. *Stem Cell Rev.* 7: 722-735.
- Kapeli, K. and Hurlin, P.J. 2011. Differential regulation of N-Myc and c-Myc synthesis, degradation, and transcriptional activity by the Ras/mitogen-activated protein kinase pathway. *J. Biol. Chem.* 286: 38498-38508.
- Regan, P.L., et al. 2011. Hsp90 inhibition increases p53 expression and destabilizes MYCN and MYC in neuroblastoma. *Int. J. Oncol.* 38: 105-112.
- Vilborg, A., et al. 2012. Wig-1, a novel regulator of N-Myc mRNA and N-Myc-driven tumor growth. *Cell Death Dis.* 3: e298.
- Masserot, C., et al. 2015. WT1 expression is inversely correlated with MYCN amplification or expression and associated with poor survival in non-MYCN-amplified neuroblastoma. *Mol. Oncol.* 10: 240-252.
- Neri, F., et al. 2015. TET1 is controlled by pluripotency-associated factors in ESCs and downmodulated by PRC2 in differentiated cells and tissues. *Nucleic Acids Res.* 43: 6814-6826.
- Fagnocchi, L., et al. 2016. A Myc-driven self-reinforcing regulatory network maintains mouse embryonic stem cell identity. *Nat. Commun.* 7: 11903.
- Kaur, G., et al. 2016. Bromodomain and hedgehog pathway targets in small cell lung cancer. *Cancer Lett.* 371: 225-239.
- Tan, J.K., et al. 2016. γ -tocotrienol acts as a BH3 mimetic to induce apoptosis in neuroblastoma SH-SY5Y cells. *J. Nutr. Biochem.* 31: 28-37.
- Qi, D.L. and Cobrinik, D. 2016. MDM2 but not MDM4 promotes retinoblastoma cell proliferation through p53-independent regulation of MYCN translation. *Oncogene* 36: 1760-1769.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA