

# TR $\alpha$ 1/ $\alpha$ 2 (2103): sc-56873

## BACKGROUND

Thyroid hormone nuclear receptors (TRs) are ligand-dependent transcription factors which regulate and control many metabolic and developmental processes. There are two genes encoding TRs identified to date, TR $\alpha$  and TR $\beta$ . TRs bind to thyroid hormone response elements (TREs) with half-site binding motifs in the orientation of palindromes, direct repeats or inverted palindromes. The affinities of binding are both variable and influenced differentially by 3,5,3'-triiodo-L-thyronine (T3). Transcriptional regulation by TRs is also modulated by heterodimerization with TR nuclear accessory proteins, the most extensively characterized of which are the retinoid X receptors (RXR $\alpha$ , RXR $\beta$  and RXR $\gamma$ ). The TR $\alpha$  isoform TR $\alpha$ 1 can display both a nuclear and undefined cytoplasmic location, and is the only TR that is imported into the mitochondrial matrix. TR $\alpha$ 2 is a C-terminal variant of TR $\alpha$ 1 that does not bind thyroid hormones (THs) and weakly binds DNA. TR $\alpha$ 2 acts as a dominant negative antagonist of TH signalling.

## REFERENCES

1. Näär, A., et al. 1991. The orientation and spacing of core DNA-binding motifs dictate selective transcriptional responses to three nuclear receptors. *Cell* 65: 1267-1271.
2. Lazar, M.A. 1993. Thyroid hormone receptors: multiple forms, multiple possibilities. *Endocrinol. Rev.* 14: 184-193.

## CHROMOSOMAL LOCATION

Genetic locus: THRA (human) mapping to 17q21.1.

## SOURCE

TR $\alpha$ 1/ $\alpha$ 2 (2103) is a mouse monoclonal antibody raised against an N-terminal peptide of TR $\alpha$ 1 of human origin.

## PRODUCT

Each vial contains IgG<sub>1</sub> in 100  $\mu$ l of 10 mM HEPES and 150 mM NaCl with < 0.1% sodium azide, 1% stabilizer protein and 25% glycerol.

## APPLICATIONS

TR $\alpha$ 1/ $\alpha$ 2 (2103) is recommended for detection of TR $\alpha$ 1/ $\alpha$ 2 of human origin by Western Blotting (starting dilution to be determined by researcher, dilution range 1:100-1:5000), immunoprecipitation [1-2  $\mu$ l per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution to be determined by researcher, dilution range 1:50-1:2500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution to be determined by researcher, dilution range 1:50-1:2500).

Suitable for use as control antibody for TR $\alpha$  siRNA (h): sc-36707, TR $\alpha$  shRNA Plasmid (h): sc-36707-SH and TR $\alpha$  shRNA (h) Lentiviral Particles: sc-36707-V.

Molecular Weight of TR $\alpha$ 1: 47 kDa.


Molecular Weight of TR $\alpha$ 2: 55 kDa.

Positive Controls: C32 nuclear extract: sc-2136, C32 whole cell lysate: sc-2205 or Hep G2 nuclear extract: sc-364819.

## STORAGE

For immediate and continuous use, store at 4° C for up to one month. For sporadic use, freeze in working aliquots in order to avoid repeated freeze/thaw cycles. If turbidity is evident upon prolonged storage, clarify solution by centrifugation.

## DATA



TR $\alpha$ 1/ $\alpha$ 2 (2103): sc-56873. Western blot analysis of TR $\alpha$ 1/ $\alpha$ 2 expression in Hep G2 nuclear extract.

## SELECT PRODUCT CITATIONS

1. Sánchez-García, O., et al. 2016. Hypothyroidism modifies morphometry and thyroid-hormone receptor expression in periurethral muscles of female rabbits. *Neurourol. Urodyn.* 35: 895-901.
2. Rodríguez-Castelán, J., et al. 2017. Distribution of thyroid hormone and thyrotropin receptors in reproductive tissues of adult female rabbits. *Endocr. Res.* 42: 59-70.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.