

IFN- α 2 (N27): sc-57219

BACKGROUND

The genes encoding type I interferons (IFNs), which include 14 IFN- α genes, one IFN- β gene, one IFN- ω (also known as IFN- α II1) gene and a number of IFN- ω pseudogenes, are clustered on human chromosome 9. Interferons- α and - β are cytokines that are widely known to induce potent antiviral activity. IFN- α and - β exert a variety of other biological effects, including antitumor and immunomodulatory activities, and are increasingly used clinically to treat a range of malignancies, myelodysplasias and autoimmune diseases. IFN- ω is antigenically different from human IFN- α , IFN- β or IFN- γ , but is a component of natural mixtures of IFN species produced by virus-induced leukocytes or Burkitt's lymphoma cells. The type I interferon receptor (IFN- α R) interacts with IFN- α , IFN- β and IFN- ω , and seems to be a multi-subunit receptor.

REFERENCES

1. Adolf, G.R. 1987. Antigenic structure of human interferon- ω 1 (interferon- α II1): comparison with other human interferons. *J. Gen. Virol.* 68: 1669-1676.
2. Lim, J.K., et al. 1994. Intrinsic ligand binding properties of the human and bovine α -interferon receptors. *FEBS Lett.* 350: 281-286.
3. Hussain, M., et al. 1996. Identification of interferon- α 7, - α 14 and - α 21 variants in the genome of a large human population. *J. Interferon Cytokine Res.* 16: 853-859.
4. Mire-Sluis, A.R., et al. 1996. An anti-cytokine bioactivity assay for interferons- α , - β and - ω . *J. Immunol. Methods* 195: 55-61.
5. Cutrone, E.C., et al. 1997. Contributions of cloned type I interferon receptor subunits to differential ligand binding. *FEBS Lett.* 404: 197-202.
6. Rozera, C., et al. 1999. Interferon (IFN)- β gene transfer into TS/A adenocarcinoma cells and comparison with IFN- α : differential effects on tumorigenicity and host response. *Am. J. Pathol.* 154: 1211-1222.
7. Barthe, C., et al. 2001. Expression of interferon- α (IFN- α) receptor 2c at diagnosis is associated with cytogenetic response in IFN- α -treated chronic myeloid leukemia. *Blood* 97: 3568-3573.
8. Eriksen, K.W., et al. 2004. Bi-phasic effect of interferon (IFN)- α : IFN- α up- and downregulates interleukin-4 signaling in human T cells. *J. Biol. Chem.* 279: 169-176.
9. Suyama, T., et al. 2005. Upregulation of the interferon- γ (IFN- γ)-inducible chemokines IFN-inducible T cell α chemoattractant and monokine induced by IFN- γ and of their receptor CXCR3 in human renal cell carcinoma. *Cancer* 103: 258-267.

CHROMOSOMAL LOCATION

Genetic locus: IFNA2 (human) mapping to 9p21.3.

SOURCE

IFN- α 2 (N27) is a mouse monoclonal antibody raised against full length IFN- α 2 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IFN- α 2 (N27) is available conjugated to either phycoerythrin (sc-57219 PE) or fluorescein (sc-57219 FITC), 200 μ g/ml, for IF, IHC(P) and FCM.

APPLICATIONS

IFN- α 2 (N27) is recommended for detection of IFN- α 2 of human origin by flow cytometry (1 μ g per 1 x 10⁶ cells); non cross-reactive with IFN- α 1.

Suitable for use as control antibody for IFN- α 2 siRNA (h): sc-63324, IFN- α 2 shRNA Plasmid (h): sc-63324-SH and IFN- α 2 shRNA (h) Lentiviral Particles: sc-63324-V.

Molecular Weight of IFN- α 2: 19 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.