

N ϵ (γ -glutamyl)-Lysine (71A3F1): sc-57600

BACKGROUND

An isopeptide bond is a unique chemical peptide bond found between a carboxyl group and an amino group in certain proteins in which the ϵ -amino groups are specified instead of the α -amino groups. This bond formed by transglutaminase is known as the N ϵ (γ -glutamyl)-Lysine bond. In this case, formation of the isopeptide bond begins with activation of ubiquitin in a process that resembles the activation of fatty acids. An adduct of ubiquitin-adenylate is formed first, followed by a thiol ester. Isopeptide linkages also occur between N- α -monomethylalanine and lysine in the ribosomal protein S11 from *Escherichia coli*.

REFERENCES

1. Goldknopf, I.L. and Busch, H. 1977. Isopeptide linkage between nonhistone and Histone 2A polypeptides of chromosomal conjugate-protein A24. *Proc. Natl. Acad. Sci. USA* 74: 864-868.
2. Chen, R. and Chen-Schmeisser, U. 1978. Isopeptide linkage between N- α -monomethylalanine and lysine in ribosomal protein S11 from *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 74: 4905-5908.
3. Kubilus, J. and Baden, H.P. 1984. Isopeptide bond formation in epidermis. *Mol. Cell. Biochem.* 58: 129-137.
4. Aswad, D.W., Johnson, B.A. and Glass, D.B. 1987. Modification of synthetic peptides related to lactate dehydrogenase (231-242) by protein carboxyl methyltransferase and tyrosine protein kinase: effects of introducing an isopeptide bond between Aspartic acid 235 and Serine 236. *Biochemistry* 26: 675-681.
5. Mezö, G., Szekerke, M., Sármay, G. and Gergely, J. 1990. Synthesis and functional studies of tuftsin analogs containing isopeptide bond. *Peptides* 11: 405-415.
6. Greco, M.A., Lorand, L., Lane, W.S., Baden, H.P., Parameswaran, K.N. and Kvedar, J.C. 1995. The pancornulins: a group of small proline rich-related cornified envelope precursors with bifunctional capabilities in isopeptide bond formation. *J. Invest. Dermatol.* 104: 204-210.
7. Friedrich, T., Kröger, B., Koerwer, W., Strube, K.H., Meyer, T. and Bialojan, S. 1998. An isopeptide bond splitting enzyme from *Hirudo medicinalis* similar to γ -glutamyl transpeptidase. *Eur. J. Biochem.* 256: 297-302.
8. Wang, P., Layfield, R., Landon, M., Mayer, R.J. and Ramage, R. 1999. Selective isopeptide bond formation: coupling ubiquitin (67-76) with Histone 2A (114-128) by use of the transfer active ester condensation technique. *J. Pept. Res.* 53: 673-677.
9. Osicka, R., Procházková, K., Sulc, M., Linhartová, I., Havlíček, V. and Sebo, P. 2004. A novel "clip-and-link" activity of repeat in toxin (RTX) proteins from Gram-negative pathogens. Covalent protein cross-linking by an Asp-Lys isopeptide bond upon calcium-dependent processing at an Asp-Pro bond. *J. Biol. Chem.* 279: 24944-24956.

SOURCE

N ϵ (γ -glutamyl)-Lysine (71A3F1) is a mouse monoclonal antibody raised against N ϵ (γ -glutamyl)-Lysine isopeptide bond.

PRODUCT

Each vial contains 50 μ g IgG2a in 500 μ l PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

N ϵ (γ -glutamyl)-Lysine (71A3F1) is recommended for detection of N ϵ (γ -glutamyl)-Lysine bond formed by transglutaminase activity by immunofluorescence (starting dilution to be determined by researcher, dilution range 1:50-1:2500); non cross-reactive with free Lysine or free glutamine; may cross-react with N ϵ (acetyl) Lysine.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.