

Giardia lamblia 65 kDa antigen (6231): sc-58131

BACKGROUND

Giardia lamblia is a flagellated protozoan parasite that infects mammalian gastrointestinal tracts and causes giardiasis, a type of gastroenteritis that manifests itself with severe diarrhea and abdominal cramps. *Giardia lamblia* infection is transmitted by consuming contaminated food or water, or through mammalian feces. The *Giardia* protozoan cannot survive in an environment outside of the host, so it is spread as a cyst, which can live in the environment for up to a month, and enters its trophozoite active form after infection. After the feeding stage, the *Giardia* trophozoite undergoes asexual replication through longitudinal binary fission. The resulting trophozoites and cysts then pass through the digestive system in the feces. The trophozoite form of *Giardia* can be recognized by its large karyosome and lack of peripheral chromatin, while the cyst form contains four nuclei and a retracted cytoplasm.

REFERENCES

1. Reiner, D.S., Douglas, H. and Gillin, F.D. 1989. Identification and localization of cyst-specific antigens of *Giardia lamblia*. *Infect. Immun.* 57: 963-968.
2. Lujan, H.D., Mowatt, M.R. and Nash, T.E. 1996. Lipid requirements and lipid uptake by *Giardia lamblia* trophozoites in culture. *J. Eukaryot. Microbiol.* 43: 237-242.
3. Hewan-Lowe, K., Furlong, B., Sims, M. and Schwartz, D.A. 1997. Coinfection with *Giardia lamblia* and *Enterocytozoon bieneusi* in a patient with acquired immunodeficiency syndrome and chronic diarrhea. *Arch. Pathol. Lab. Med.* 121: 417-422.
4. Svård, S.G., Hagblom, P. and Palm, J.E. 2003. *Giardia lamblia*—a model organism for eukaryotic cell differentiation. *FEMS Microbiol. Lett.* 218: 3-7.
5. Benchimol, M. 2004. Participation of the adhesive disc during karyokinesis in *Giardia lamblia*. *Biol. Cell* 96: 291-301.
6. Benchimol, M. 2004. *Giardia lamblia*: behavior of the nuclear envelope. *Parasitol. Res.* 94: 254-264.
7. Chávez-Munguía, B., Cedillo-Rivera, R. and Martínez-Palomo, A. 2004. The ultrastructure of the cyst wall of *Giardia lamblia*. *J. Eukaryot. Microbiol.* 51: 220-226.
8. Corrêa, G. and Benchimol, M. 2006. *Giardia lamblia* behavior under cytochalasins treatment. *Parasitol. Res.* 98: 250-256.
9. Misra, V., Misra, S.P., Dwivedi, M. and Singh, P.A. 2006. *Giardia lamblia* trophozoites in gastric biopsies. *Indian J. Pathol. Microbiol.* 49: 519-523.

SOURCE

Giardia lamblia 65 kDa antigen (6231) is a mouse monoclonal antibody raised against *Giardia lamblia*.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Giardia lamblia 65 kDa antigen (6231) is recommended for detection of the 65 kDa antigen of *Giardia lamblia* by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of Giardia lamblia 65 kDa antigen: 65 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.