
SANTA CRUZ BIOTECHNOLOGY, INC.

ABCG2 (BXP-53): sc-58224



BACKGROUND

ATP-binding cassette (ABC) transporters are an evolutionarily conserved family of proteins that catalyze the transport of molecules across extracellular and intracellular membranes through the energy of ATP hydrolysis. The ABC half-transporter, ABCG2, is also known as placenta-specific ABC transporter and breast cancer resistance protein (BCRP1). ABCG2 confers resistance for a variety of chemotherapeutic agents, including anthracyclines, mitoxantrone, bisantrene and topotecan. Under normal conditions, ABCG2 may serve a protective function by removing toxins from the cell, and plays an important role in regulating stem cell differentiation. ABCG2 is responsible for the side population (SP) phenotype and is widely expressed in a large variety of stem cells, making it an important stem cell marker. ABCG2 may have N-linked glycosylation and may dimerize *in vivo*. ABCG2 is abundantly expressed in placenta, liver, intestine and stem cells.

CHROMOSOMAL LOCATION

Genetic locus: ABCG2 (human) mapping to 4q22.1; Abcg2 (mouse) mapping to 6 B3.

SOURCE

ABCG2 (BXP-53) is a rat monoclonal antibody raised against amino acids 221-394 of ABCG2 of mouse origin.

PRODUCT

Each vial contains 100 μg lgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% stabilizer protein.

APPLICATIONS

ABCG2 (BXP-53) is recommended for detection of ABCG2 of mouse, rat and human origin by Western Blotting (starting dilution to be determined by researcher, dilution range 1:10-1:200), immunoprecipitation [10-20 μ l per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution to be determined by researcher, dilution range 1:10-1:200) and flow cytometry (10-20 μ l per 1 x 10⁶ cells).

Suitable for use as control antibody for ABCG2 siRNA (h): sc-41151, ABCG2 siRNA (m): sc-37054, ABCG2 shRNA Plasmid (h): sc-41151-SH, ABCG2 shRNA Plasmid (m): sc-37054-SH, ABCG2 shRNA (h) Lentiviral Particles: sc-41151-V and ABCG2 shRNA (m) Lentiviral Particles: sc-37054-V.

Molecular Weight of ABCG2: 72 kDa.

Positive Controls: ABCG2 (h3): 293T Lysate: sc-172804, JAR cell lysate: sc-2276 or MCF7 whole cell lysate: sc-2206.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


PROTOCOLS


See our web site at www.scbt.com for detailed protocols and support products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA





ABCG2 (BXP-53): sc-58224. Western blot analysis of ABCG2 expression in HL-60/MX-1 (A), JAR (B) and MCF7 (C) whole cell lysates.

ABCG2 (BXP-53): sc-58224. Western blot analysis of ABCG2 expression in non-transfected: sc-117752 (A) and human ABCG2 transfected: sc-172804 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- 1. Solazzo, M., et al. 2009. Mitochondrial expression and functional activity of breast cancer resistance protein in different multiple drug-resistant cell lines. Cancer Res. 69: 7235-7242.
- 2. Guo, S., et al. 2014. Higher level of plasma bioactive molecule sphingosine 1-phosphate in women is associated with estrogen. Biochim. Biophys. Acta 1841: 836-846.
- Bhagwandin, V.J., et al. 2016. The metastatic potential and chemoresistance of human pancreatic cancer stem cells. PLoS ONE 11: e0148807.
- Nickel, S., et al. 2017. Expression and activity of breast cancer resistance protein (BCRP/ABCG2) in human distal lung epithelial cells *in vitro*. Pharm. Res. 34: 2477-2487.
- Lamartinière, Y., et al. 2018. ABCA7 downregulation modifies cellular cholesterol homeostasis and decreases amyloid-β peptide efflux in an *in vitro* model of the blood-brain barrier. J. Alzheimers Dis. 64: 1195-1211.
- Yellamilli, A., et al. 2020. Abcg2-expressing side population cells contribute to cardiomyocyte renewal through fusion. FASEB J. 34: 5642-5657.
- Tatebayashi, R., et al. 2021. Gene-expression profile and postpartum transition of bovine endometrial side population cells0. Biol. Reprod. 104: 850-860.
- Shin, J., et al. 2022. Endothelial OCT4 is atheroprotective by preventing metabolic and phenotypic dysfunction. Cardiovasc. Res. 118: 2458-2477.
- Gunes, Y., et al. 2023. Modulation of monepantel secretion into milk by soy isoflavones. J. Vet. Pharmacol. Ther. 46: 185-194.



See **ABCG2 (B-1): sc-377176** for ABCG2 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.