

CD4 (YTS 191.1.2): sc-59042

BACKGROUND

The T cell receptor (TCR) is a heterodimer composed of either α and β or γ and δ chains. CD3 chains and the CD4 or CD8 co-receptors are also required for efficient signal transduction through the TCR. The TCR is expressed on T helper and T cytotoxic cells that can be distinguished by their expression of CD4 and CD8; T helper cells express CD4 proteins and T cytotoxic cells display CD8. CD4 is also expressed on cortical cells, mature medullary thymocytes, microglial cells and Dendritic cells. CD4 (also designated T4 and Leu 3), is a 55 kDa membrane glycoprotein that contains four extracellular immunoglobulin-like domains. The TCR in association with CD4 can bind class II MHC molecules presented by the antigen-presenting cells. The CD4 protein functions by increasing the avidity of the interaction between the TCR and an antigen-class II MHC complex. An additional role of CD4 is to function as a receptor for HIV.

REFERENCES

- Maddon, P.J., et al. 1987. Structure and expression of human and mouse T4 genes. Proc. Natl. Acad. Sci. USA 84: 9155-9159.
- Arthos, J., et al. 1989. Identification of the residues in human CD4 critical for the binding of HIV. Cell 57: 469-481.
- Healey, D., et al. 1990. Novel anti-CD4 monoclonal antibodies separate human immunodeficiency virus infection and fusion of CD4⁺ cells from virus binding. J. Exp. Med. 172: 1233-1242.
- Allison, J.P., et al. 1991. The immunobiology of T cells with invariant γ δ antigen receptors. Annu. Rev. Immunol. 9: 679-705.
- Janeway, C.A., Jr. 1992. The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation. Annu. Rev. Immunol. 10: 645-674.
- Ehrlich, E.W., et al. 1993. T cell receptor interaction with peptide/major histocompatibility complex (MHC) and superantigen/MHC ligands is dominated by antigen. J. Exp. Med. 178: 713-722.
- Julius, M., et al. 1993. Distinct roles for CD4 and CD8 as coreceptors in antigen receptor signalling. Immunol. Today 14: 177-183.
- Vignali, D.A. 1994. The interaction between CD4 and MHC class II molecules and its effect on T cell function. Behring Inst. Mitt. 94: 133-147.

CHROMOSOMAL LOCATION

Genetic locus: Cd4 (mouse) mapping to 6 F2.

SOURCE

CD4 (YTS 191.1.2) is a rat monoclonal antibody raised against CD4 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CD4 (YTS 191.1.2) is recommended for detection of CD4 of mouse origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1×10^6 cells).

Suitable for use as control antibody for CD4 siRNA (m): sc-29997, CD4 shRNA Plasmid (m): sc-29997-SH and CD4 shRNA (m) Lentiviral Particles: sc-29997-V.

Molecular Weight of CD4: 54 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **CD4 (MT310): sc-19641** for CD4 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.