

MPO (MPO-7): sc-59604

BACKGROUND

The heme protein myeloperoxidase (MPO) is a major component of azurophilic granules of neutrophils and polymorphonuclear leukocytes. Optimal oxygen-dependent microbiocidal activity depends on MPO as the critical enzyme for the generation of hypochlorous acid and other toxic oxygen products. The MPO precursor is synthesized during the promyelocytic stage of myeloid differentiation and is subsequently processed and transported intracellularly to the lysosomes. The precursor undergoes cotranslational N-linked glycosylation to produce a glycoprotein. Glucosidases in the endoplasmic reticulum (ER) or early *cis* Golgi convert the pro-MPO to a form which is sorted into a prelysosomal compartment, which undergoes final proteolytic maturation to native MPO, a pair of heavy-light protomers. In normal neutrophils, MPO is expressed as a dimer. Calreticulin, a calcium-binding protein residing in the ER, interacts specifically with fully glycosylated apopro-MPO. iMPO mRNA is abundant in human promyelocytic HL-60 and mouse myeloid leukemia NFS-60 cells. MPO is expressed at high levels in circulating neutrophils and monocytes but is not detectable in microglia, brain-specific macrophages or normal brain tissue.

REFERENCES

1. Johnson, K.R., Nauseef, W.M., Care, A., Wheelock, M.J., Shane, S., Hudson, S., Koeffler, H.P., Selsted, M., Miller, C. and Rovera, G. 1987. Characterization of cDNA clones for human myeloperoxidase: predicted amino acid sequence and evidence for multiple mRNA species. *Nucleic Acids Res.* 15: 2013-2028.
2. Nauseef, W.M. 1987. Posttranslational processing of a human myeloid lysosomal protein, myeloperoxidase. *Blood* 70: 1143-1150.
3. Morishita, K., Kubota, N., Asano, S., Kazi, Y. and Nagata, S. 1987. Molecular cloning and characterization of cDNA for human myeloperoxidase. *J. Biol. Chem.* 262: 3844-3851.
4. Nauseef, W.M., Olsson, I. and Amljots, K. 1988. Biosynthesis and processing of myeloperoxidase a marker for myeloid cell differentiation. *Eur. J. Haematol.* 40: 97-110.
5. Homma, T., Suzuki, K., Kudo, Y., Inagawa, M., Mizuno, S., Yamaguchi, K. and Tagawa, M. 1989. Preparation and characterization of monoclonal antibodies against human myeloperoxidase. *Arch. Biochem. Biophys.* 273: 189-196.
6. Zurbier, K.W., van den Berg, J.D., Van Gelder, B.F. and Muijsers, A.O. 1992. Human hemi-myeloperoxidase. Initial chlorinating activity at neutral pH, compound II and III formation, and stability towards hypochlorous acid and high temperature. *Eur. J. Biochem.* 205: 737-742.
7. Nauseef, W.M., McCormick, S.J. and Clark, R.A. 1995. Calreticulin functions as a molecular chaperone in the biosynthesis of myeloperoxidase. *J. Biol. Chem.* 270: 4741-4747.
8. Reynolds, W.F., Rhees, J., Maciejewski, D., Paladino, T., Sieburg, H., Maki, R.A. and Masliah, E. 1999. Myeloperoxidase polymorphism is associated with gender specific risk for Alzheimer's disease. *Exp. Neurol.* 155: 31-41.

CHROMOSOMAL LOCATION

Genetic locus: MPO (human) mapping to 17q23.1.

SOURCE

MPO (MPO-7) is a mouse monoclonal antibody raised against MPO from CML cells of human origin.

PRODUCT

Each vial contains 250 µl culture supernatant containing IgG₁ with < 0.1% sodium azide.

APPLICATIONS

MPO (MPO-7) is recommended for detection of MPO of human origin by immunofluorescence (starting dilution to be determined by researcher, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution to be determined by researcher, dilution range 1:50-1:500).

Suitable for use as control antibody for MPO siRNA (h): sc-43941, MPO shRNA Plasmid (h): sc-43941-SH and MPO shRNA (h) Lentiviral Particles: sc-43941-V.

Molecular Weight of native MPO heavy-light protomer: 72 kDa.

Molecular Weight of MPO dimer: 140 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.