AP-3 δ siRNA (h): sc-60176



The Power to Question

BACKGROUND

Clathrin-coated pits and vesicles are assembled for receptor-mediated endocytosis through interaction with Clathrin associated protein complexes. Vesicle transport is mediated from the *trans*-Golgi network by the adapter complex AP-1 and from the plasma membrane by the AP-2 complex. AP-3 (also designated AP180 or F1-20) is a synapse-specific Clathrin assembly protein. The protein CALM (Clathrin assembly protein lymphoid myeloid leukemia) is highly homologous to AP180 and may also be involved in Clathrin assembly. AP-3 α , AP-3 α and AP-3 α are important parts of the AP-3 complex.

REFERENCES

- 1. Robinson, M.S. 1989. Cloning of cDNAs encoding two related 100-kD coated vesicle proteins (α -Adaptins). J. Cell Biol. 108: 833-842.
- 2. Kirchhausen, T., et al. 1989. Structural and functional division into two domains of the large (100- to 115-kDa) chains of the clathrin-associated protein complex AP-2. Proc. Natl. Acad. Sci. USA 86: 2612-2616.
- Robinson, M.S. 1990. Cloning and expression of γ-Adaptin, a component of clathrin-coated vesicles associated with the Golgi apparatus. J. Cell Biol. 111: 2319-2326.
- Ponnambalam, S., et al. 1990. Conservation and diversity in families of coated vesicle adaptins. J. Biol. Chem. 265: 4814-4820.
- Simpson, F., et al. 1997. Characterization of the adaptor-related protein complex, AP-3. J. Cell Biol. 137: 835-845.
- Lefrancois, S., et al. 2004. An ear-core interaction regulates the recruitment of the AP-3 complex to membranes. Dev. Cell 7: 619-625.

CHROMOSOMAL LOCATION

Genetic locus: AP3D1 (human) mapping to 19p13.3.

PRODUCT

AP-3 δ siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AP-3 δ shRNA Plasmid (h): sc-60176-SH and AP-3 δ shRNA (h) Lentiviral Particles: sc-60176-V as alternate gene silencing products.

For independent verification of AP-3 δ (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60176A, sc-60176B and sc-60176C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AP-3 δ siRNA (h) is recommended for the inhibition of AP-3 δ expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AP-3 δ gene expression knockdown using RT-PCR Primer: AP-3 δ (h)-PR: sc-60176-PR (20 μ I). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

 Subramanian, G., et al. 2024. AP3B1 has type I interferon-independent antiviral function against SARS-CoV-2. Viruses 16: 1377.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Furope +00800 4573 8000 49 6221 4503 0 www.scbt.com