

BBS8 siRNA (h): sc-60261

BACKGROUND

Bardet-Biedl syndrome (BBS) is a heterogeneous pleiotropic genetic disorder characterized by obesity, photoreceptor degeneration, polydactyly, hypogonadism, renal abnormalities, developmental delay, diabetes, hypertension and congenital heart defects. BBS genes map to eight genetic loci and encode eight proteins, BBS1-BBS8. Five BBS genes encode basal body or cilia proteins, suggesting that BBS is a ciliary dysfunction disorder. Mutations in BBS8, also designated tetratricopeptide repeat protein (TTC8), probably account for only a minority (2%) of BBS families, underlining the difficulty of genotyping heterogeneous conditions. The identification of BBS8 provides the key to the pathogenesis of the condition as a primary ciliary disorder.

REFERENCES

1. Young, J.C., et al. 1998. Specific binding of tetratricopeptide repeat proteins to the C-terminal 12-kDa domain of hsp90. *J. Biol. Chem.* 273: 18007-18010.
2. Cortajarena, A.L., et al. 2004. Protein design to understand peptide ligand recognition by tetratricopeptide repeat proteins. *Protein Eng. Des. Sel.* 17: 399-409.
3. Heon, E., et al. 2005. Ocular phenotypes of three genetic variants of Bardet-Biedl syndrome. *Am. J. Med. Genet. A* 132A: 283-287.
4. Nakane, T., et al. 2005. No evidence for triallelic inheritance of MKKS/BBS loci in Amish McKusick-Kaufman syndrome. *Am. J. Med. Genet. A* 138: 32-34.
5. Hichri, H., et al. 2005. Testing for triallelism: analysis of six BBS genes in a Bardet-Biedl syndrome family cohort. *Eur. J. Hum. Genet.* 13: 607-616.
6. Dollfus, H., et al. 2005. Update on Bardet-Biedl syndrome. *J. Fr. Ophtalmol.* 28: 106-112.
7. Nachury, M.V., et al. 2007. A core complex of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane biogenesis. *Cell* 129: 1201-1213.

CHROMOSOMAL LOCATION

Genetic locus: TTC8 (human) mapping to 14q31.3.

PRODUCT

BBS8 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see BBS8 shRNA Plasmid (h): sc-60261-SH and BBS8 shRNA (h) Lentiviral Particles: sc-60261-V as alternate gene silencing products.

For independent verification of BBS8 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60261A, sc-60261B and sc-60261C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

BBS8 siRNA (h) is recommended for the inhibition of BBS8 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

BBS8 (E-2): sc-271009 is recommended as a control antibody for monitoring of BBS8 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor BBS8 gene expression knockdown using RT-PCR Primer: BBS8 (h)-PR: sc-60261-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.