

# C6ST-1 siRNA (m): sc-60306

## BACKGROUND

Sulfotransferase enzymes catalyze the sulfate conjugation of many hormones, neurotransmitters, drugs and xenobiotic compounds. These cytosolic enzymes differ in their tissue distribution and substrate specificities, although the gene structure (number and length of exons) is similar among family members. Sulfotransferases are primarily expressed in liver and adrenal tissues, where they add sulfate to steroids and bile acids. Chondroitin 6-sulfotransferase-1 (C6ST-1) is a 486 amino acid protein that localizes in the Golgi apparatus, where it sulfates both chondroitin and keratan sulfate. C6ST-1 is developmentally regulated in many different tissues, with expression continuing through adulthood in the spleen. When C6ST-1 expression is upregulated, the motility of Schwann cells that guide growing axons through both developmental and injured environments increases.

## REFERENCES

1. Fukuta, M., et al. 1995. Molecular cloning and expression of chick chondrocyte chondroitin 6-sulfotransferase. *J. Biol. Chem.* 270: 18575-18580.
2. Gauguet, J.M., et al. 2004. Core 2 branching  $\beta$ 1-6-N-acetylglucosaminyltransferase and high endothelial cell N-acetylglucosamine-6-sulfotransferase exert differential control over B and T lymphocyte homing to peripheral lymph nodes. *Blood* 104: 4104-4112.
3. Uchimura, K., et al. 2004. N-acetylglucosamine-6-O-sulfotransferase-1 regulates expression of L-Selectin ligands and lymphocyte homing. *J. Biol. Chem.* 279: 35001-35008.
4. de Graffenried, C.L. and Bertozzi, C.R. 2004. The stem region of the sulfotransferase GlcNAc6ST-1 is a determinant of substrate specificity. *J. Biol. Chem.* 279: 40035-40043.
5. Thiele, H., et al. 2004. Loss of chondroitin 6-O-sulfotransferase-1 function results in severe human chondrodysplasia with progressive spinal involvement. *Proc. Natl. Acad. Sci. USA* 101: 10155-10160.
6. Yamada, T., et al. 2004. Chondroitin 4-sulphotransferase-1 and chondroitin 6-sulphotransferase-1 are affected differently by uronic acid residues neighboring the acceptor GalNAc residues. *Biochem. J.* 384: 567-575.

## CHROMOSOMAL LOCATION

Genetic locus: Chst3 (mouse) mapping to 10 B4.

## PRODUCT

C6ST-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see C6ST-1 shRNA Plasmid (m): sc-60306-SH and C6ST-1 shRNA (m) Lentiviral Particles: sc-60306-V as alternate gene silencing products.

For independent verification of C6ST-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60306A, sc-60306B and sc-60306C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

C6ST-1 siRNA (m) is recommended for the inhibition of C6ST-1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

C6ST-1 (G-9): sc-271696 is recommended as a control antibody for monitoring of C6ST-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor C6ST-1 gene expression knockdown using RT-PCR Primer: C6ST-1 (m)-PR: sc-60306-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.