CHMP2 siRNA (h): sc-60369



The Power to Question

BACKGROUND

The charged multivesicular body proteins, commonly designated CHMPs, belong to the vacuolar sorting protein family and function as chromatin-modifying proteins. CHMP1-6 are all components of ESCRT (endosomal sorting complex required for transport) I, II or III complexes. These complexes are crucial for sorting endosomal articles into multivesicular bodies (MVBs), and are also required for the formation of these bodies. CHMP2, also known as BC-2, associates directly with Vps4 for the disassembly of ESCRT-III complex in an ATP-dependant manner. During HIV-1 infection, the virus uses the ESCRT-III complex to mediate budding and exocytosis of viral proteins. Overexpression of CHMP2 strongly inhibits HIV-1 release.

REFERENCES

- von Schwedler, U.K., Stuchell, M., Müller, B., Ward, D.M., Chung, H.Y., Morita, E., Wang, H.E., Davis, T., He, G.P., Cimbora, D.M., Scott, A., Kräusslich, H.G., Kaplan, J., Morham, S.G. and Sundquist, W.I. 2003. The protein network of HIV budding. Cell 114: 701-713.
- 2. Suo, Y.P., Wang, B.Y., Hong, Z., Yin, R.T., Wang, D.Q. and Peng, Z.L. 2006. Comparative study of that synergistic effect of cisplatin combined with compound herbal medicinal prescription for tonic quality and activating blood circulation is on SKOV3 cell proliferation and apoptosis. Sichuan Da Xue Xue Bao Yi Xue Ban 37: 542-546.
- Agromayor, M. and Martin-Serrano, J. 2006. Interaction of AMSH with ESCRT-III and deubiquitination of endosomal cargo. J. Biol. Chem. 281: 23083-23091.

CHROMOSOMAL LOCATION

Genetic locus: CHMP2A (human) mapping to 19q13.43.

PRODUCT

CHMP2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CHMP2 shRNA Plasmid (h): sc-60369-SH and CHMP2 shRNA (h) Lentiviral Particles: sc-60369-V as alternate gene silencing products.

For independent verification of CHMP2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60369A, sc-60369B and sc-60369C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CHMP2 siRNA (h) is recommended for the inhibition of CHMP2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CHMP2 gene expression knockdown using RT-PCR Primer: CHMP2 (h)-PR: sc-60369-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

 Hemmati, F., Akinpelu, A., Song, J., Amiri, F., McDaniel, A., McMurray, C., Afthinos, A., Andreadis, S.T., Aitken, A.V., Biancardi, V.C., Gerecht, S. and Mistriotis, P. 2023. Downregulation of YAP activity restricts P53 hyperactivation to promote cell survival in confinement. Adv. Sci. 10: e2302228.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com