

# CRHSP-24 siRNA (h): sc-60443

## BACKGROUND

CRHSP-24 (calcium-regulated heat-stable protein 24) is a serine phosphoprotein originally identified as a physiological substrate for the  $\text{Ca}^{2+}$ -calmodulin regulated protein phosphatase calcineurin (PP2B). A ubiquitously expressed protein, CRHSP-24 interacts with the STYX/dead phosphate protein in developing spermatids and is a paralog of the brain-specific mRNA-binding protein PIPPIN. It is thought to play a common role in calcium-mediated signal transduction resulting from phosphorylation on serine residues. CRHSP-24 is phosphorylated on Ser 52 by Akt-1 (PKB $\alpha$ ) in response to IGF-1, on Ser 52 by Akt-1 and RSK in response to EGF, and on Ser 41 in the absence of IGF-1/EGF by a DYRK isoform or another proline-directed protein kinase.

## REFERENCES

1. Burnham, D.B., et al. 1982. Effects of carbachol, cholecystokinin, and insulin on protein phosphorylation in isolated pancreatic acini. *J. Biol. Chem.* 257: 10523-10528.
2. Goblewski, G.E., et al. 1998. Purification and characterization of a novel physiological substrate for calcineurin in mammalian cells. *J. Biol. Chem.* 273: 22738-22744.
3. Williams, J.A., et al. 1999. Intracellular regulatory mechanisms in pancreatic acinar cellular function. *Curr. Opin. Gastroenterol.* 15: 385-391.
4. Williams, J.A., et al. 2001. Intracellular signaling mechanisms activated by cholecystokinin-regulating synthesis and secretion of digestive enzymes in pancreatic acinar cells. *Annu. Rev. Physiol.* 63: 77-97.
5. Wishart, M.J., et al. 2002. PTEN and myotubularin phosphatases: from 3-phosphoinositide dephosphorylation to disease. *Trends Cell Biol.* 12: 579-585.
6. Schäfer, C., et al. 2003. CRHSP-24 phosphorylation is regulated by multiple signaling pathways in acinar cells. *Am. J. Physiol. Gastrointest. Liver Physiol.* 285: G726-G734.
7. Sans, M.D., et al. 2004. Calcineurin is required for translational control of protein synthesis in rat pancreatic acini. *Am. J. Physiol., Cell Physiol.* 287: C310-C319.

## CHROMOSOMAL LOCATION

Genetic locus: CRHSP1 (human) mapping to 16p13.2.

## PRODUCT

CRHSP-24 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CRHSP-24 shRNA Plasmid (h): sc-60443-SH and CRHSP-24 shRNA (h) Lentiviral Particles: sc-60443-V as alternate gene silencing products.

For independent verification of CRHSP-24 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60443A, sc-60443B and sc-60443C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}\text{C}$ , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

CRHSP-24 siRNA (h) is recommended for the inhibition of CRHSP-24 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

CRHSP-24 (A-1): sc-137072 is recommended as a control antibody for monitoring of CRHSP-24 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CRHSP-24 gene expression knockdown using RT-PCR Primer: CRHSP-24 (h)-PR: sc-60443-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^{\circ}\text{C}$  and the extension temperature should be  $68-72^{\circ}\text{C}$ .

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.