



ENT3 siRNA (h): sc-60587

BACKGROUND

Equilibrative nucleoside transporters (ENTs) regulate many physiological processes and are widely distributed in mammals, plants, yeasts, insects, nematodes and protozoans. They enable facilitated diffusion of hydrophilic nucleosides, such as adenosine, and nucleoside analogs across cell membranes. ENTs are required for uptake of antiviral and anticancer nucleoside drugs and influence a variety of physiological processes, such as neurotransmission and platelet aggregation, by regulating the amount of adenoside available to cell surface receptors. Equilibrative nucleoside transporter 3 (ENT3), also designated solute carrier family 29 (nucleoside transporters), member 3, belongs to the SLC29A transporter family and is a mammalian ENT isoform. ENT3 functions intracellularly, rather than at the plasma membrane, and may export nucleosides from the lysosomal interior.

REFERENCES

- Hyde, R.J., et al. 2001. The ENT family of eukaryote nucleoside and nucleobase transporters: recent advances in the investigation of structure/function relationships and the identification of novel isoforms. *Mol. Membr. Biol.* 18: 53-63.
- Alcorn, J., et al. 2002. Transporter gene expression in lactating and non-lactating human mammary epithelial cells using real-time reverse transcription-polymerase chain reaction. *J. Pharmacol. Exp. Ther.* 303: 487-496.
- Acimovic, Y. and Coe, I.R. 2002. Molecular evolution of the equilibrative nucleoside transporter family: identification of novel family members in prokaryotes and eukaryotes. *Mol. Biol. Evol.* 19: 2199-2210.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: Q9BZD2. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Lu, H., et al. 2004. Tissue distribution of concentrative and equilibrative nucleoside transporters in male and female rats and mice. *Drug Metab. Dispos.* 32: 1455-1461.
- Baldwin, S.A., et al. 2005. Functional characterization of novel human and mouse equilibrative nucleoside transporters (hENT3 and mENT3) located in intracellular membranes. *J. Biol. Chem.* 280: 15880-15887.

CHROMOSOMAL LOCATION

Genetic locus: SLC29A3 (human) mapping to 10q22.1.

PRODUCT

ENT3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ENT3 shRNA Plasmid (h): sc-60587-SH and ENT3 shRNA (h) Lentiviral Particles: sc-60587-V as alternate gene silencing products.

For independent verification of ENT3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60587A, sc-60587B and sc-60587C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ENT3 siRNA (h) is recommended for the inhibition of ENT3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ENT3 gene expression knockdown using RT-PCR Primer: ENT3 (h)-PR: sc-60587-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.