

IL-27R α siRNA (m): sc-60837

BACKGROUND

IL-27 is a heterodimeric cytokine that consists of EBI3, an IL-12p40-related protein and p28, a IL-12p35-related polypeptide. IL-27 triggers expansion of antigen-specific naive CD4⁺ T cells and promotes polarization towards a Th1 phenotype with expression of γ -interferon. IL-27 contributes to the development of an adaptive immune response through its action on CD4⁺ T cells, and also directly acts on cells of the innate immune system. IL-27 protein levels increase upon activation of antigen-presenting cells. IL-27 protein induces orphan cytokine receptor IL-27R (WSX-1)-dependent clonal expansion of naive but not memory CD4⁺ T cells. IL-27 signaling through IL-27R and gp130 also induces phosphorylation of Stat1-5.

REFERENCES

1. Pflanz, S., et al. 2002. IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4⁺ T cells. *Immunity* 16: 779-790.
2. Cordoba-Rodriguez, R. and Frucht, D.M. 2003. IL-23 and IL-27: new members of the growing family of IL-12-related cytokines with important implications for therapeutics. *Expert Opin. Biol. Ther.* 3: 715-723.
3. Lucas, S., et al. 2003. IL-27 regulates IL-12 responsiveness of naive CD4⁺ T cells through Stat1-dependent and -independent mechanisms. *Proc. Natl. Acad. Sci. USA* 100: 15047-15052.
4. Villarino, A.V., et al. 2004. Understanding the pro- and anti-inflammatory properties of IL-27. *J. Immunol.* 173: 715-720.
5. Goldberg, R., et al. 2004. Suppression of ongoing adjuvant-induced arthritis by neutralizing the function of the p28 subunit of IL-27. *J. Immunol.* 173: 1171-1178.
6. Yoshimoto, T., et al. 2004. Induction of IgG_{2a} class switching in B cells by IL-27. *J. Immunol.* 173: 2479-2485.
7. Artis, D., et al. 2004. The IL-27 receptor (WSX-1) is an inhibitor of innate and adaptive elements of type 2 immunity. *J. Immunol.* 173: 5626-5634.
8. Holscher, C., et al. 2005. The IL-27 receptor chain WSX-1 differentially regulates antibacterial immunity and survival during experimental tuberculosis. *J. Immunol.* 174: 3534-3544.

CHROMOSOMAL LOCATION

Genetic locus: Il27ra (mouse) mapping to 8 C3.

PRODUCT

IL-27R α siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-27R α shRNA Plasmid (m): sc-60837-SH and IL-27R α shRNA (m) Lentiviral Particles: sc-60837-V as alternate gene silencing products.

For independent verification of IL-27R α (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60837A, sc-60837B and sc-60837C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IL-27R α siRNA (m) is recommended for the inhibition of IL-27R α expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-27R α gene expression knockdown using RT-PCR Primer: IL-27R α (m)-PR: sc-60837-PR (20 μ l, 598 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.