



# Lipin-3 siRNA (m): sc-60945

## BACKGROUND

The Lipin family of nuclear proteins contains three members: Lipin-1, Lipin-2, and Lipin-3, all of which contain a nuclear signal sequence, a highly conserved amino-terminal (NLIP) domain, and a carboxy-terminal (CLIP) domain. Lipin-1 is crucial for normal adipose tissue development and metabolism. Lipin-1 selectively activates a subset of PGC-1 $\alpha$  target pathways, including fatty acid oxidation and mitochondrial oxidative phosphorylation by inducing expression of the nuclear receptor PPAR $\alpha$ . Lipin-1 also inactivates the lipogenic program and suppresses circulating lipid levels. Lipin-2 is linked to Majeed syndrome, and autosomal recessive, autoinflammatory disorder. Lipin-3 is an 851 amino acid protein that localizes to the nucleus. Lipin-3 observations are useful in studies related to adipose tissue development in the context of obesity, fatty liver dystrophy, lipodystrophy, Insulin resistance and Type 2 diabetes.

## REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605520. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Phan, J., Peterfy, M. and Reue, K. 2004. Lipin expression preceding peroxisome is critical for adipogenesis *in vivo* and *in vitro*. J. Biol. Chem. 279: 29558-29564.
3. Reitman, M.L. 2005. The fat and thin of Lipin. Cell Metab. 1: 5-6.
4. Phan, J. and Reue, K. 2005. Lipin, a lipodystrophy and obesity gene. Cell Metab. 1: 73-83.
5. Scavallo, G.S., Paluru, P.C., Zhou, J., White, P.S., Rappaport, E.F. and Young, T.L. 2005. Genomic structure and organization of the high grade Myopia-2 locus (MYP2) critical region: mutation screening of 9 positional candidate genes. Mol. Vis. 11: 97-110.
6. Han, G.S., Wu, W.I. and Carman, G.M. 2006. The *Saccharomyces cerevisiae* Lipin homolog is a Mg<sup>2+</sup>-dependent phosphatidate phosphatase enzyme. J. Biol. Chem. 281: 9210-9218.
7. Parsons, T.R. 2006. Studies on Lipin-protein complexes: lecithin-caseinogen complexes. Biochem. J. 22: 800-810.
8. Suviolahti, E., Reue, K., Cantor, R.M., Phan, J., Gentile, M., Naukkarinen, J., Soro-Paavonen, A., Oksanen, L., Kaprio, J., Rissanen, A., Salomaa, V., Kontula, K., Taskinen, M.R., Pajukanta, P. and Peltonen, L. 2006. Cross-species analyses implicate Lipin-1 involvement in human glucose metabolism. Hum. Mol. Genet. 15: 377-386.

## CHROMOSOMAL LOCATION

Genetic locus: Lpin3 (mouse) mapping to 2 H2.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

Lipin-3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Lipin-3 shRNA Plasmid (m): sc-60945-SH and Lipin-3 shRNA (m) Lentiviral Particles: sc-60945-V as alternate gene silencing products.

For independent verification of Lipin-3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60945A, sc-60945B and sc-60945C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Lipin-3 siRNA (m) is recommended for the inhibition of Lipin-3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Lipin-3 gene expression knockdown using RT-PCR Primer: Lipin-3 (m)-PR: sc-60945-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.