

Lubricin siRNA (m): sc-60973

BACKGROUND

Lubricin, also designated proteoglycan-4 or megakaryocyte stimulating factor, plays an important role in boundary lubrication within articulating joints. The disulfide-linked dimer of Lubricin, bonded between Cys 1146 and Cys 1403, is essential for protein cleavage. Highly expressed in cartilage, liver and synovial tissue, Lubricin inhibits synovial cell adhesion to the cartilage surface, but also prevents the deposition of proteins from synovial fluid into cartilage. Defects in the gene encoding for Lubricin can cause Jakobs syndrome, also designated campptodactyly-arthropathy-coxa vara-pericarditis syndrome (CACP). CACP is an autosomal recessive disorder characterized by joint failure associated with noninflammatory synoviocyte hyperplasia and subintimal fibrosis of the synovial capsule. Lubricin undergoes different levels of glycosylation and may be detected at varying molecular weights.

REFERENCES

1. Schaefer, D.B., et al. 2004. Lubricin reduces cartilage—cartilage integration. *Biorheology* 41: 503-508.
2. Elsaid, K.A., et al. 2005. Association of articular cartilage degradation and loss of boundary-lubricating ability of synovial fluid following injury and inflammatory arthritis. *Arthritis Rheum.* 52: 1746-1755.
3. Kontinen, Y.T., et al. 2005. The microenvironment around total hip replacement prostheses. *Clin. Orthop. Relat. Res.* 430: 28-38.
4. Rhee, D.K., et al. 2005. Consequences of disease-causing mutations on Lubricin protein synthesis, secretion, and post-translational processing. *J. Biol. Chem.* 280: 31325-31332.

CHROMOSOMAL LOCATION

Genetic locus: Prg4 (mouse) mapping to 1 G1.

PRODUCT

Lubricin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Lubricin shRNA Plasmid (m): sc-60973-SH and Lubricin shRNA (m) Lentiviral Particles: sc-60973-V as alternate gene silencing products.

For independent verification of Lubricin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60973A, sc-60973B and sc-60973C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Lubricin siRNA (m) is recommended for the inhibition of Lubricin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Lubricin gene expression knockdown using RT-PCR Primer: Lubricin (m)-PR: sc-60973-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Liu, S.Q., et al. 2012. Cardioprotective proteins upregulated in the liver in response to experimental myocardial ischemia. *Am. J. Physiol. Heart Circ. Physiol.* 303: H1446-H1458.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.