



# NALP11 siRNA (h): sc-61142

## BACKGROUND

The NACHT-, LRR- and PYD-containing protein (NALP) family functions in the regulation of apoptosis and inflammatory signaling pathways. Members of the NALP family (also designated Pyrin-containing APAF1-like proteins) include NALP1 through NALP11. Several family members, such as NALP1, NALP2, NALP3 and NALP6, influence NF $\kappa$ B and caspase pathways as components of the inflammasome. NALP11 is a 1,033-amino acid member of the NALP family with the typical PYD-NACHT-LRR domain structure. It is implicated in the activation of proinflammatory caspases via its involvement in multiprotein complexes called inflammasomes.

## REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 609662. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Chamaillard, M., et al. 2003. NODS, NALPS and NAIP: intracellular regulators of bacterial-induced inflammation. *Cell. Microbiol.* 5: 581-592.
3. Tschopp, J., et al. 2003. NALPs: a novel protein family involved in inflammation. *Nat. Rev. Mol. Cell Biol.* 4: 95-104.
4. Hiller, S., et al. 2003. NMR structure of the apoptosis- and inflammation-related NALP1 Pyrin domain. *Structure* 11: 1199-1205.
5. Damiano, J.S., et al. 2004. Heterotypic interactions among NACHT domains: implications for regulation of innate immune responses. *Biochem. J.* 381: 213-219.
6. Sanz, C., et al. 2004. NALP1 is a transcriptional target for cAMP-response-element-binding protein (CREB) in myeloid leukaemia cells. *Biochem. J.* 384: 281-286.
7. Liu, F., et al. 2004. Expression of NALP1 in cerebellar granule neurons stimulates apoptosis. *Cell. Signal.* 16: 1013-1021.
8. Kinoshita, T., et al. 2005. PYPAF3, a PYRIN-containing APAF-1-like protein, is a feedback regulator of caspase-1-dependent interleukin-1 $\beta$  secretion. *J. Biol. Chem.* 280: 21720-21725.

## CHROMOSOMAL LOCATION

Genetic locus: NLRP11 (human) mapping to 19q13.42.

## PRODUCT

NALP11 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NALP11 shRNA Plasmid (h): sc-61142-SH and NALP11 shRNA (h) Lentiviral Particles: sc-61142-V as alternate gene silencing products.

For independent verification of NALP11 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61142A, sc-61142B and sc-61142C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

NALP11 siRNA (h) is recommended for the inhibition of NALP11 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NALP11 gene expression knockdown using RT-PCR Primer: NALP11 (h)-PR: sc-61142-PR (20  $\mu$ l, 546 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Gangopadhyay, A., et al. 2022. NLRP3 licenses NLRP11 for inflammasome activation in human macrophages. *Nat. Immunol.* 23: 892-903.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.