

OATP-E siRNA (m): sc-61248

BACKGROUND

The organic anion transporter family of proteins mediate hepatic uptake of cardiac glycosides. OATP-E (organic anion transporter E), also known as SLC04A1 (solute carrier organic anion transporter family member 4A1), SLC21A12 (solute carrier family 21 member 12) or POAT, is a 722 amino acid member of the organic anion transporter protein family. As a multi-pass membrane protein, OATP-E mediates the Na⁺-independent transport of estrone-3-sulfate, taurocholate and the thyroid hormones T3 (triiodo-L-thyronine), T4 (thyroxine) and rT3. OATP-E is ubiquitously expressed except in leukocytes and spleen. OATP-E is expressed as four isoforms produced by alternative splicing events.

REFERENCES

1. Tamai, I., Nezu, J., Uchino, H., Sai, Y., Oku, A., Shimane, M. and Tsuji, A. 2000. Molecular identification and characterization of novel members of the human organic anion transporter (OATP) family. *Biochem. Biophys. Res. Commun.* 273: 251-260.
2. Fujiwara, K., Adachi, H., Nishio, T., Unno, M., Tokui, T., Okabe, M., Onogawa, T., Suzuki, T., Asano, N., Tanemoto, M., Seki, M., Shiiba, K., Suzuki, M., Kondo, Y., et al. 2001. Identification of thyroid hormone transporters in humans: different molecules are involved in a tissue-specific manner. *Endocrinology* 142: 2005-2012.
3. Ito, A., Yamaguchi, K., Tomita, H., Suzuki, T., Onogawa, T., Sato, T., Mizutamari, H., Mikkaichi, T., Nishio, T., Suzuki, T., Unno, M., Sasano, H., Abe, T. and Tamai, M. 2003 Distribution of rat organic anion transporting polypeptide-E (Oatp-E) in the rat eye. *Invest. Ophthalmol. Vis. Sci.* 44: 4877-4884.
4. Sato, K., Sugawara, J., Sato, T., Mizutamari, H., Suzuki, T., Ito, A., Mikkaichi, T., Onogawa, T., Tanemoto, M., Unno, M., Abe, T. and Okamura, K. 2003. Expression of organic anion transporting polypeptide E (Oatp-E) in human placenta. *Placenta* 24: 144-148.
5. Nozawa, T., Suzuki, M., Takahashi, K., Yabuuchi, H., Maeda, T., Tsuji, A. and Tamai, I. 2004. Involvement of estrone-3-sulfate transporters in proliferation of hormone-dependent breast cancer cells. *J. Pharmacol. Exp. Ther.* 311: 1032-1037.

CHROMOSOMAL LOCATION

Genetic locus: Slco4a1 (mouse) mapping to 2 H4.

PRODUCT

OATP-E siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see OATP-E shRNA Plasmid (m): sc-61248-SH and OATP-E shRNA (m) Lentiviral Particles: sc-61248-V as alternate gene silencing products.

For independent verification of OATP-E (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61248A, sc-61248B and sc-61248C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

OATP-E siRNA (m) is recommended for the inhibition of OATP-E expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor OATP-E gene expression knockdown using RT-PCR Primer: OATP-E (m)-PR: sc-61248-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.