

Rev-erb β siRNA (h): sc-61456

BACKGROUND

Orphan nuclear receptors NR1D1 and NR1D2 are more commonly designated Rev-erb α and Rev-erb β , respectively. Rev-erb α acts as a receptor for tri-iodothyronine and is composed of three domains: a modulating N-terminal domain, a C-terminal steroid binding domain and a DNA-binding domain. Rev-erb β binds to the sequences 5'-AATGTAGGTCA-3' and 5'-ATAACTAGGTCA-3' and acts as a competitive repressor of ROR α function. It interacts with NCOA5 co-activator which leads to an increase in transcription. Both Rev-erb α and Rev-erb β are nuclear proteins belonging to the nuclear hormone receptor family of proteins.

REFERENCES

1. Laudet, V., Begue, A., Henry-Duthoit, C., Joubel, A., Martin, P., Stehelin, D. and Saule, S. 1991. Genomic organization of the human thyroid hormone receptor α (c-erbA-1) gene. *Nucleic Acids Res.* 19: 1105-1112.
2. Dumas, B., Harding, H.P., Choi, H.S., Lehmann, K.A., Chung, M., Lazar, M.A. and Moore, D.D. 1995. A new orphan member of the nuclear hormone receptor superfamily closely related to Rev-erb. *Mol. Endocrinol.* 8: 996-1005.
3. Kainu, T., Enmark, E., Gustafsson, J.A. and Peltö-Huikko, M.P. 1996. Localization of the Rev-erbA orphan receptors in the brain. *Brain Res.* 743: 315-319.
4. Zhao, Q., Khorasanizadeh, S., Miyoshi, Y., Lazar, M.A. and Rastinejad, F. 1998. Structural elements of an orphan nuclear receptor-DNA complex. *Mol. Cell* 1: 849-861.
5. Koh, Y.S. and Moore, D.D. 1999. Linkage of the nuclear hormone receptor genes NR1D2, THRB and RARB: evidence for an ancient, large-scale duplication. *Genomics* 57: 289-292.
6. Sauve, F., McBroom, L.D., Gallant, J., Moraitis, A.N., Labrie, F. and Giguère, V. 2001. CIA, a novel estrogen receptor coactivator with a bifunctional nuclear receptor interacting determinant. *Mol. Cell. Biol.* 21: 343-353.
7. Kamphuis, W., Cailotto, C., Dijk, F., Bergen, A. and Buijs, R.M. 2005. Circadian expression of clock genes and clock-controlled genes in the rat retina. *Biochem. Biophys. Res. Commun.* 330: 18-26.

CHROMOSOMAL LOCATION

Genetic locus: NR1D2 (human) mapping to 3p24.2.

PRODUCT

Rev-erb β siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rev-erb β shRNA Plasmid (h): sc-61456-SH and Rev-erb β shRNA (h) Lentiviral Particles: sc-61456-V as alternate gene silencing products.

For independent verification of Rev-erb β (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61456A, sc-61456B and sc-61456C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rev-erb β siRNA (h) is recommended for the inhibition of Rev-erb β expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Rev-erb β (D-8): sc-398252 is recommended as a control antibody for monitoring of Rev-erb β gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rev-erb β gene expression knockdown using RT-PCR Primer: Rev-erb β (h)-PR: sc-61456-PR (20 μ l, 539 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.