

SDF-4 siRNA (m): sc-61509

BACKGROUND

The C-X-C or α chemokine family is characterized by a pair of cysteine residues separated by a single amino acid and primarily functions as chemoattractants for neutrophils. The C-X-C family includes IL-8, NAP-2, MSGA, stromal cell derived factor-1 (SDF-1) and SDF-4. SDF-1 was originally described as a pre-B cell stimulatory factor, but has now been shown to function as a potent chemoattractant for T cells and monocytes but not neutrophils. SDF-4, also designated Cab45, contains six EF-hand calcium-binding motifs. SDF-4 is ubiquitously expressed and localizes to the Golgi where it may regulate calcium-dependent activities.

REFERENCES

1. Deng, H., Liu, R., Ellmeier, W., Choe, S., Unutmaz, D., Burkhart, M., Di Marzio, P., Marmon, S., Sutton, R.E. and Hill, C.M. 1996. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 381: 661-666.
2. Nagasawa, T., Hirota, S., Tachibana, K., Takakura, N., Nishikawa, S., Kitamura, Y., Yoshida, N., Kikutani, H. and Kishimoto, T. 1996. Defects of B cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature* 382: 635-638.
3. Bleul, C.C., Farzan, M., Choe, H., Parolin, C., Clark-Lewis, I., Sodroski, J. and Springer, T.A. 1996. The lymphocyte chemoattractant SDF-1 is a ligand for LESTR/fusin and blocks HIV-1 entry. *Nature* 382: 829-833.
4. Laterveer, L., Lindley, I.J., Heemskerk, D.P., Camps, J.A., Pauwels, E.K., Willemze, R. and Fibbe, W.E. 1996. Rapid mobilization of hematopoietic progenitor cells in Rhesus monkeys by a single intravenous injection of interleukin-8. *Blood* 87: 781-788.
5. Scherer, P.E., Lederkremer, G.Z., Williams, S., Fogliano, M., Baldini, G. and Lodish, H.F. 1996. Cab45, a novel Ca^{2+} -binding protein localized to the Golgi lumen. *J. Cell Biol.* 133: 257-268.
6. Koivu, T., Laitinen, S., Riento, K. and Olkkonen, V.M. 1997. Sequence of a human cDNA encoding Cab45, a Ca^{2+} -binding protein with six EF-hand motifs. *DNA Seq.* 7: 217-220.

CHROMOSOMAL LOCATION

Genetic locus: Sdf4 (mouse) mapping to 4 E2.

PRODUCT

SDF-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SDF-4 shRNA Plasmid (m): sc-61509-SH and SDF-4 shRNA (m) Lentiviral Particles: sc-61509-V as alternate gene silencing products.

For independent verification of SDF-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61509A, sc-61509B and sc-61509C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SDF-4 siRNA (m) is recommended for the inhibition of SDF-4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

SDF-4 (E-12): sc-393930 is recommended as a control antibody for monitoring of SDF-4 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SDF-4 gene expression knockdown using RT-PCR Primer: SDF-4 (m)-PR: sc-61509-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.