



## TaR siRNA (h): sc-61644

### BACKGROUND

Trace amines are endogenous molecules structurally related to classical biogenic amines that are linked to psychiatric conditions. A family of G protein-coupled receptors referred to as trace-amine-associated receptors, (TaRs or TAARs) are activated by trace amines and are present in very low levels in mammalian tissue. TaRs contain several structural features that are similar to the rhodopsin  $\beta$ -adrenergic receptor superfamily, including the positions of the seven transmembrane regions that provide common ligand-binding pockets as well as the short N- and C-terminal domains. TaRs are associated with the detection of social cues, illustrating their significance as therapeutic targets. Specifically, TaR proteins are potential targets for drugs of abuse, such as amphetamine and MDMA, as well as neuropsychiatric disorders including schizophrenia, depression, and attention deficit disorder.

### REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604849. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Aridon, P., D'Andrea, G., Rigamonti, A., Leone, M., Casari, G. and Bussone, G. 2004. Elusive amines and cluster headache: mutational analysis of trace amine receptor cluster on chromosome 6q23. *Neurol. Sci.* 25: S279-S280.
3. Lindemann, L., Ebeling, M., Kratochwil, N.A., Bunzow, J.R., Grandy, D.K. and Hoener, M.C. 2005. Trace amine-associated receptors form structurally and functionally distinct subfamilies of novel G protein-coupled receptors. *Genomics* 85: 372-385.
4. Miller, G.M., Verrico, C.D., Jassen, A., Konar, M., Yang, H., Panas, H., Bahn, M., Johnson, R. and Madras, B.K. 2005. Primate trace amine receptor 1 modulation by the dopamine transporter. *J. Pharm. Exp. Ther.* 313: 983-994.
5. Liberles, S.D. and Buck, L.B. 2006. A second class of chemosensory receptors in the olfactory epithelium. *Nature* 442: 645-650.
6. Wainscott, D.B., Little, S.P., Yin, T., Tu, Y., Rocco, V.P., He, J.X. and Nelson, D.L. 2006. Pharmacologic characterization of the cloned human trace amine-associated receptor1 (TAAR1) and evidence for species differences with the rat TAAR1. *J. Pharmacol. Exp. Ther.* 320: 475-485.

### CHROMOSOMAL LOCATION

Genetic locus: TAAR1 (human) mapping to 6q23.2.

### PRODUCT

TaR siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TaR shRNA Plasmid (h): sc-61644-SH and TaR shRNA (h) Lentiviral Particles: sc-61644-V as alternate gene silencing products.

For independent verification of TaR (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61644A, sc-61644B and sc-61644C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

TaR siRNA (h) is recommended for the inhibition of TaR expression in human cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TaR gene expression knockdown using RT-PCR Primer: TaR (h)-PR: sc-61644-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.