



Grx5 siRNA (h): sc-61857

BACKGROUND

Glutaredoxin (Grx) and a relative, thioredoxin, catalyze general thiol-disulfide oxidoreductions and act as hydrogen donors for ribonucleotide reductase, an enzyme essential for DNA synthesis. Proteins which catalyze thiol-disulfide exchange reactions are required for electron and proton transport to essential enzymes like ribonucleotide reductase, for the formation of disulfide bonds during protein folding, and for general regulation of protein function by thiol redox control. Grx proteins also play a role in cellular defense against oxidative stress. The thioredoxin superfamily includes a number of proteins with the same basic folding and structure as thioredoxin and glutaredoxin, with the active site at the C-terminal end of a β -strand followed by an α -helix. Glutaredoxin (Grx) operates in thiol-disulfide reactions via two vicinal (CXYC) active site cysteine residues, which either form a disulfide (oxidized form) or a dithiol (reduced form). Grx-5 (Glutaredoxin-related protein 5), also known as GLRX5, is a 157 amino acid mitochondrial glutaredoxin necessary for the function of iron/sulfur enzymes in yeast.

REFERENCES

1. Björnstedt, M., et al. 1997. Selenium and the thioredoxin and glutaredoxin systems. *Biomed. Environ. Sci.* 10: 271-279.
2. Rodríguez-Manzanique, M.T., et al. 2002. Grx5 is a mitochondrial glutaredoxin required for the activity of iron/sulfur enzymes. *Mol. Biol. Cell* 13: 1109-1121.
3. Fernandes, A.P. and Holmgren, A. 2004. Glutaredoxins: glutathione-dependent redox enzymes with functions far beyond a simple thioredoxin backup system. *Antioxid. Redox Signal.* 6: 63-74.
4. Vilella, F., et al. 2004. Evolution and cellular function of monothiol glutaredoxins: involvement in iron-sulphur cluster assembly. *Comp. Funct. Genomics* 5: 328-341.
5. Alves, R., et al. 2004. Predictive reconstruction of the mitochondrial iron-sulfur cluster assembly metabolism. II. Role of glutaredoxin Grx5. *Proteins* 57: 481-492.
6. Wingert, R.A., et al. 2005. Deficiency of glutaredoxin 5 reveals Fe-S clusters are required for vertebrate haem synthesis. *Nature* 436: 1035-1039.
7. Herrero, E., et al. 2006. Glutaredoxins in fungi. *Photosyn. Res.* 89: 127-140.
8. Herrero, E. and de la Torre-Ruiz, M.A. 2007. Monothiol glutaredoxins: a common domain for multiple functions. *Cell. Mol. Life Sci.* 64: 1518-1530.
9. Kalinina, E.V., et al. 2008. Involvement of thio-, peroxi-, and glutaredoxins in cellular redox-dependent processes. *Biochemistry* 73: 1493-1510.

CHROMOSOMAL LOCATION

Genetic locus: GLRX5 (human) mapping to 14q32.13.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Grx5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Grx5 shRNA Plasmid (h): sc-61857-SH and Grx5 shRNA (h) Lentiviral Particles: sc-61857-V as alternate gene silencing products.

For independent verification of Grx5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61857A, sc-61857B and sc-61857C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Grx5 siRNA (h) is recommended for the inhibition of Grx5 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Grx5 gene expression knockdown using RT-PCR Primer: Grx5 (h)-PR: sc-61857-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.