

Tau siRNA (r): sc-61900

BACKGROUND

Tau can be phosphorylated by several protein kinases. Phosphorylation by Tau protein kinase II at Serine 404 is primarily responsible for the functional loss of Tau-mediated tubulin polymerization. In addition, phosphorylation of microtubule-associated Tau results in the dissociation of Tau from the microtubules and tubulin depolymerization. Serine 412 of Tau is modified by Tau protein kinase I/glycogen synthase kinase-3 β (TPKI/GSK-3 β) to disrupt neuronal metabolism in anatomical areas vulnerable to Alzheimer's disease. TPKI/GSK-3 β is expressed primarily in neurons and especially in neurites early in development, whereafter the distribution is concentrated mostly in the cell soma and the proximal neurite region.

REFERENCES

1. Cross, D., et al. 1993. A Tau-like protein interacts with stress fibers and microtubules in human and rodent cultured cell lines. *J. Cell Sci.* 105: 51-60.
2. Lubke, U., et al. 1994. Microtubule-associated protein Tau epitopes are present in fiber lesions in diverse muscle disorders. *Am. J. Pathol.* 145: 175-188.
3. Singh, T.J., et al. 1996. Differential phosphorylation of human Tau isoforms containing three repeats by several protein kinases. *Arch. Biochem. Biophys.* 328: 43-50.
4. Hoshi, M., et al. 1996. Regulation of mitochondrial pyruvate dehydrogenase activity by Tau protein kinase I/glycogen synthase kinase-3 β in brain. *Proc. Natl. Acad. Sci. USA* 93: 2719-2723.
5. Malchiodi-Albedi, F., et al. 1997. Protein phosphatase inhibitors induce modification of synapse structure and Tau hyperphosphorylation in cultured rat hippocampal neurons. *J. Neurosci. Res.* 48: 425-438.

CHROMOSOMAL LOCATION

Genetic locus: Mapt (rat) mapping to 10q32.1.

PRODUCT

Tau siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Tau shRNA Plasmid (r): sc-61900-SH and Tau shRNA (r) Lentiviral Particles: sc-61900-V as alternate gene silencing products.

For independent verification of Tau (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61900A, sc-61900B and sc-61900C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Tau siRNA (r) is recommended for the inhibition of Tau expression in rat cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Tau (Tau 46): sc-32274 is recommended as a control antibody for monitoring of Tau gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Tau gene expression knockdown using RT-PCR Primer: Tau (r)-PR: sc-61900-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.