

# Astrotactin 2 siRNA (h): sc-62000

## BACKGROUND

Neuronal migration of the central nervous system is a specialized form of cell motility that takes place in the laminar structure of the cortical regions of brain. Astrotactin is a neuronal cell surface protein expressed on postmitotic neuronal precursors in the cerebellum, hippocampus, cerebrum, and olfactory bulb. Astrotactin 2 is a paralog of Astrotactin. Astrotactin mediates neuron-astroglial interactions and is also implicated in synaptic development as well as many other neuronal activities. Astrotactin has three epidermal growth factor repeat domains and two fibronectin type III repeat domains. The human Astrotactin gene shows extensive homology to the mouse Astrotactin gene. Mutations in the Astrotactin gene are linked to neuronal migration defects in both species.

## REFERENCES

1. Edmondson, J.C., et al. 1988. Astrotactin: a novel neuronal cell surface antigen that mediates neuron-astroglial interactions in cerebellar microcultures. *J. Cell Biol.* 106: 505-517.
2. Stitt, T.N., et al. 1990. Antibodies that recognize Astrotactin block granule neuron binding to astroglia. *Neuron* 5: 639-649.
3. Fishell, G., et al. 1992. Astrotactin provides a receptor system for CNS neuronal migration. *Development* 113: 755-765.
4. Zheng, C., et al. 1996. CNS gene encoding Astrotactin, which supports neuronal migration along glial fibers. *Science* 272: 417-419.
5. Adams, N.C., et al. 2002. Mice that lack Astrotactin have slowed neuronal migration. *Development* 129: 965-972.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 600904. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Chen, Y.T., et al. 2005. Deficits in motor coordination with aberrant cerebellar development in mice lacking testicular orphan nuclear receptor 4. *Mol. Cell. Biol.* 25: 2722-2732.
8. Tárnok, K., et al. 2005. Cerebellar granule cells show age-dependent migratory differences *in vitro*. *J. Neurobiol.* 65: 135-145.

## CHROMOSOMAL LOCATION

Genetic locus: ASTN2 (human) mapping to 9q33.1.

## PRODUCT

Astrotactin 2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Astrotactin 2 shRNA Plasmid (h): sc-62000-SH and Astrotactin 2 shRNA (h) Lentiviral Particles: sc-62000-V as alternate gene silencing products.

For independent verification of Astrotactin 2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62000A, sc-62000B and sc-62000C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Astrotactin 2 siRNA (h) is recommended for the inhibition of Astrotactin 2 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Astrotactin 2 gene expression knockdown using RT-PCR Primer: Astrotactin 2 (h)-PR: sc-62000-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.