



COX7c siRNA (m): sc-62153

BACKGROUND

The cytochrome c oxidase (COX) family of proteins function as the final electron donor in the respiratory chain to drive a proton gradient across the inner mitochondrial membrane, ultimately resulting in the production of water. The mammalian COX apoenzyme is a dimer, with each monomer consisting of 13 subunits, some of which are mitochondrial and some of which are nuclear. The COX7c protein is a nuclear-encoded subunit of the COX complex that is transported to the mitochondria where it functions with the other COX proteins in the last step in cellular respiration. Like most of the other cytochrome subunits, COX7c is aerobic and its expression is increased when oxygen levels are high.

REFERENCES

1. Seelan, R.S. and Grossman, L.I. 1997. Structural organization and promoter analysis of the bovine cytochrome c oxidase subunit VIIc gene. A functional role for YY1. *J. Biol. Chem.* 272: 10175-10181.
2. Burke, P.V., Raitt, D.C., Allen, L.A., Kellogg, E.A. and Poyton, R.O. 1997. Effects of oxygen concentration on the expression of cytochrome c and cytochrome c oxidase genes in yeast. *J. Biol. Chem.* 272: 14705-14712.
3. Ragno, S., Estrada-Garcia, I., Butler, R. and Colston, M.J. 1998. Regulation of macrophage gene expression by *Mycobacterium tuberculosis*: down-regulation of mitochondrial cytochrome c oxidase. *Infect. Immun.* 66: 3952-3958.
4. Schmidt, T.R., Goodman, M. and Grossman, L.I. 2002. Amino acid replacement is rapid in primates for the mature polypeptides of COX subunits, but not for their targeting presequences. *Gene* 286: 13-19.
5. Yu, L., Lei, L.S. and Wu, S.G. 2003. Cyclosporin A inhibits Insulin release and down-regulates gene expressions of mitochondrial oxidative phosphorylation enzymes in NIT-cells. *Di Yi Jun Yi Da Xue Xue Bao* 23: 833-836.
6. De Angelis, P.M., Svendsrud, D.H., Kravik, K.L. and Stokke, T. 2006. Cellular response to 5-fluorouracil (5-FU) in 5-FU-resistant colon cancer cell lines during treatment and recovery. *Mol. Cancer* 5: 20.
7. Dang, C., Gottschling, M., Manning, K., O'Curraín, E., Schneider, S., Sterry, W., Stockfleth, E. and Nindl, I. 2006. Identification of dysregulated genes in cutaneous squamous cell carcinoma. *Oncol. Rep.* 16: 513-519.

CHROMOSOMAL LOCATION

Genetic locus: Cox7c (mouse) mapping to 13 C3.

PRODUCT

COX7c siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see COX7c shRNA Plasmid (m): sc-62153-SH and COX7c shRNA (m) Lentiviral Particles: sc-62153-V as alternate gene silencing products.

For independent verification of COX7c (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62153A, sc-62153B and sc-62153C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

COX7c siRNA (m) is recommended for the inhibition of COX7c expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.