



CTL4 siRNA (h): sc-62167

BACKGROUND

Choline is an essential nutrient that is required for the synthesis of both acetylcholine, a neurotransmitter found in cholinergic nerve terminals, and phosphatidylcholine, a key component of cell membranes. Choline deficiencies are associated with defects in cell growth and have been implicated in disorders such as Alzheimer's and Parkinson's disease. The choline transporter-like protein family (CTL) are solute carriers that transport choline, a compound which is not able to permeate cells, across the cell membrane. CTL4, also known as SLC44A4 (solute carrier family 44 member 4), is a multi-pass membrane protein which can fuse with Neu1, generating a CTL4-Neu1 transcript. This fusion is implicated in sialidosis, a disease characterized by improper lysosomal storage.

REFERENCES

1. Uhl, J., et al. 2002. Identification of a CTL4/Neu1 fusion transcript in a sialidosis patient. *FEBS Lett.* 521: 19-23.
2. Traiffort, E., et al. 2005. Molecular characterization of the family of choline transporter-like proteins and their splice variants. *J. Neurochem.* 92: 1116-1125.
3. Michel, V., et al. 2006. Choline transport for phospholipid synthesis. *Exp. Biol. Med.* 231: 490-504.
4. Sjöblom, T., et al. 2006. The consensus coding sequences of human breast and colorectal cancers. *Science* 314: 268-274.
5. Tomi, M., et al. 2007. Na⁺-independent choline transport in rat retinal capillary endothelial cells. *Neurochem. Res.* 32: 1833-1842.
6. Wang, T., et al. 2007. Choline transporters in human lung adenocarcinoma: expression and functional implications. *Acta Biochim. Biophys. Sin.* 39: 668-674.

CHROMOSOMAL LOCATION

Genetic locus: SLC44A4 (human) mapping to 6p21.33.

PRODUCT

CTL4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CTL4 shRNA Plasmid (h): sc-62167-SH and CTL4 shRNA (h) Lentiviral Particles: sc-62167-V as alternate gene silencing products.

For independent verification of CTL4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62167A, sc-62167B and sc-62167C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CTL4 siRNA (h) is recommended for the inhibition of CTL4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CTL4 gene expression knockdown using RT-PCR Primer: CTL4 (h)-PR: sc-62167-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.