

DPRP1 siRNA (m): sc-62235

BACKGROUND

Dipeptidyl peptidases (DPPs) mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. DPRP1 (dipeptidyl-peptidase IV-related protein 1), also known as DPP8 (dipeptidyl-peptidase 8), DP8 or MSTP141, is a member of the peptidase S9B family of proteins that exhibit prolyl oligopeptidase activity. DPRP1 localizes to the cytoplasm and is ubiquitously expressed with predominant expression in placenta, brain, prostate, testis and muscle. DPRP1 is similar to CD26 (dipeptidyl peptidase IV) suggesting that it may be involved in immune function and participate in the activation of T-cells.

REFERENCES

1. Abbott, C.A., et al. 2000. Cloning, expression and chromosomal localization of a novel human dipeptidyl peptidase (DPP) IV homolog, DPP8. *Eur. J. Biochem.* 267: 6140-6150.
2. Olsen, C., et al. 2002. Identification and characterization of human DPP9, a novel homologue of dipeptidyl peptidase IV. *Gene* 299: 185-193.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606819. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Qi, S.Y., et al. 2003. Cloning and characterization of dipeptidyl peptidase 10, a new member of an emerging subgroup of serine proteases. *Biochem. J.* 373: 179-189.
5. Busek, P., et al. 2007. Dipeptidyl peptidase-IV enzymatic activity bearing molecules in human brain tumors—good or evil? *Front. Biosci.* 13: 2319-2326.

CHROMOSOMAL LOCATION

Genetic locus: Dpp8 (mouse) mapping to 9 C.

PRODUCT

DPRP1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DPRP1 shRNA Plasmid (m): sc-62235-SH and DPRP1 shRNA (m) Lentiviral Particles: sc-62235-V as alternate gene silencing products.

For independent verification of DPRP1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62235A, sc-62235B and sc-62235C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DPRP1 siRNA (m) is recommended for the inhibition of DPRP1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

DPRP1 (A-5): sc-376399 is recommended as a control antibody for monitoring of DPRP1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DPRP1 gene expression knockdown using RT-PCR Primer: DPRP1 (m)-PR: sc-62235-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.