



H3-K9-HMTase siRNA (m): sc-62430

BACKGROUND

H3-K9-HMTase (histone-lysine N-methyltransferase SETDB2) is a 719 amino acid protein encoded by the human gene SETDB2. H3-K9-HMTase, which belongs to the histone-lysine methyltransferase family, contains one MBD (methyl-CpG-binding) domain, one pre-SET domain and one SET domain. H3-K9-HMTase is believed to be a probable histone methyltransferase with catalytic activity. Epigenetic gene silencing in eukaryotes is regulated in part by lysine methylation of the core histone proteins. While histone lysine methylation is known to control gene expression through the recruitment of modification-specific effector proteins, it remains unknown whether non-histone chromatin proteins are targets for similar modification-recognition systems. Located in the nucleus, H3-K9-HMTase is ubiquitously expressed with highest expression found in heart, testis and ovary.

REFERENCES

1. Ebbs, M.L., Bartee, L. and Bender, J. 2005. H3 lysine 9 methylation is maintained on a transcribed inverted repeat by combined action of SUVH6 and SUVH4 methyltransferases. *Mol. Cell. Biol.* 25: 10507-10515.
2. Estève, P.O., Patnaik, D., Chin, H.G., Benner, J., Teitell, M.A. and Pradhan, S. 2005. Functional analysis of the N- and C-terminus of mammalian G9a histone H3 methyltransferase. *Nucleic Acids Res.* 33: 3211-3223.
3. Tachibana, M., Ueda, J., Fukuda, M., Takeda, N., Ohta, T., Iwanari, H., Sakihama, T., Kodama, T., Hamakubo, T. and Shinkai, Y. 2005. Histone methyltransferases G9a and GLP form heteromeric complexes and are both crucial for methylation of euchromatin at H3-K9. *Genes Dev.* 19: 815-826.
4. Nishijima, H., Nakayama, J., Yoshioka, T., Kusano, A., Nishitani, H., Shibahara, K. and Nishimoto, T. 2006. Nuclear RanGAP is required for the heterochromatin assembly and is reciprocally regulated by histone H3 and Ctr4 histone methyltransferase in *Schizosaccharomyces pombe*. *Mol. Biol. Cell* 17: 2524-2536.
5. Tyagi, S., Chabes, A.L., Wysocka, J. and Herr, W. 2007. E2F activation of S phase promoters via association with HCF-1 and the MLL family of histone H3K4 methyltransferases. *Mol. Cell* 27: 107-119.
6. Henkels, C.H. and Khorasanizadeh, S. 2007. Implications of a histone code mimic in epigenetic signaling. *Mol. Cell* 27: 521-522.
7. Tzeng, T.Y., Lee, C.H., Chan, L.W. and Shen, C.K. 2007. Epigenetic regulation of the Drosophila chromosome 4 by the histone H3K9 methyltransferase dSETDB1. *Proc. Natl. Acad. Sci. USA* 104: 12691-12696.

CHROMOSOMAL LOCATION

Genetic locus: Setdb2 (mouse) mapping to 14 C3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

H3-K9-HMTase siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see H3-K9-HMTase shRNA Plasmid (m): sc-62430-SH and H3-K9-HMTase shRNA (m) Lentiviral Particles: sc-62430-V as alternate gene silencing products.

For independent verification of H3-K9-HMTase (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62430A, sc-62430B and sc-62430C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

H3-K9-HMTase siRNA (m) is recommended for the inhibition of H3-K9-HMTase expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor H3-K9-HMTase gene expression knockdown using RT-PCR Primer: H3-K9-HMTase (m)-PR: sc-62430-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.