



Malcavernin siRNA (h): sc-62594

BACKGROUND

Cerebral cavernous malformation (CCM) is an autosomal dominant or sporadic neurovascular disease marked by vascular anomalies located mostly in the central nervous system that can cause stroke, seizures, cerebral hemorrhages, headaches and focal neurologic deficits. CCM is caused by mutations in one of three genes: CCM1, CCM2 or CCM3. CCM1 encodes the protein KRIT1, CCM2 encodes the protein Malcavernin and CCM3 shares its name with the protein it encodes. Malcavernin, also designated cerebral cavernous malformations 2 protein, is a scaffolding protein for MEK kinase-3. Like KRIT1, Malcavernin is expressed in a variety of human organs including the arterial vascular endothelium, pyramidal neurons, astrocytes and their foot processes. In addition, Malcavernin is expressed in various epithelial cells that are required for the formation of the blood-organ barrier. Malcavernin is localized to the cytoplasm but is known to shuttle to and from the nucleus. Due to its lack of a nuclear export signal or nuclear localization signal, it is believed that Malcavernin accomplishes this shuttling via an attachment to KRIT1, which contains a nuclear localization signal. Two isoforms exist for Malcavernin. Isoform 1 represents the full length protein while isoform 2 contains an alternative four amino acid sequence rather than the first 10 residues of isoform 1.

REFERENCES

1. Lawton, M.T., et al. 2005. Giant infiltrative cavernous malformation: clinical presentation, intervention, and genetic analysis: case report. *Neurosurgery* 55: 979-980.
2. Guclu, B., et al. 2005. Mutations in apoptosis-related gene, PDCD10, cause cerebral cavernous malformation 3. *Neurosurgery* 57: 1008-1013.
3. Guclu, B., et al. 2005. Cerebral venous malformations have distinct genetic origin from cerebral cavernous malformations. *Stroke* 36: 2479-2480.
4. Zawistowski, J.S., et al. 2005. CCM1 and CCM2 protein interactions in cell signaling: implications for cerebral cavernous malformations pathogenesis. *Hum. Mol. Genet.* 14: 2521-2531.

CHROMOSOMAL LOCATION

Genetic locus: CCM2 (human) mapping to 7p13.

PRODUCT

Malcavernin siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Malcavernin shRNA Plasmid (h): sc-62594-SH and Malcavernin shRNA (h) Lentiviral Particles: sc-62594-V as alternate gene silencing products.

For independent verification of Malcavernin (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62594A, sc-62594B and sc-62594C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Malcavernin siRNA (h) is recommended for the inhibition of Malcavernin expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Malcavernin gene expression knockdown using RT-PCR Primer: Malcavernin (h)-PR: sc-62594-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.