

# Munc13-4 siRNA (m): sc-62652

## BACKGROUND

Munc13-4 is a member of the Munc13 family and is a homolog to Munc13-1. Munc13-4 lacks the C1 domain and N-terminal extension that are present in other Munc13 family members. It is a peripheral membrane, GTP-Rab 27a-binding protein. Munc13-4 has a ubiquitous tissue distribution; however, unlike related proteins Munc13-1, -2 and -3, Munc13-4 is mainly expressed outside the nervous system. High expression levels of Munc13-4 have been seen in mucous goblet and alveolar type II cells of the lung, as well as in cytotoxic T lymphocytes and mast cells. Munc13-4 localizes to secretory lysosomes. Overexpression of Munc13-4 enhances degranulation of mast cell secretory lysosomes, suggesting that it positively regulates secretory lysosome fusion and exocytosis. Mutations in Munc13-4 cause familial hemophagocytic lymphohistiocytosis subtype 3.

## REFERENCES

1. Feldmann, J., et al. 2003. Munc13-4 is essential for cytolytic granules fusion and is mutated in a form of familial hemophagocytic lymphohistiocytosis (FHL-3). *Cell* 115: 461-473.
2. Shirakawa, R., et al. 2004. Munc13-4 is a GTP-Rab 27-binding protein regulating dense core granule secretion in platelets. *J. Biol. Chem.* 279: 10730-10737.
3. Neef, M., et al. 2005. Munc13-4 is an effector of Rab 27a and controls secretion of lysosomes in hematopoietic cells. *Mol. Biol. Cell* 16: 731-741.
4. Ishii, E., et al. 2005. Genetic subtypes of familial hemophagocytic lymphohistiocytosis: correlations with clinical features and cytotoxic T lymphocyte/natural killer cell functions. *Blood* 105: 3442-3448.
5. Yamamoto, K., et al. 2005. Mutations of Syntaxin 11 and SNAP 23 genes as causes of familial hemophagocytic lymphohistiocytosis were not found in Japanese people. *J. Hum. Genet.* 50: 600-603.
6. Hong, W., et al. 2005. Cytotoxic T lymphocyte exocytosis: bring on the SNAREs! *Trends Cell Biol.* 15: 644-650.
7. Schneider, E.M., et al. 2006. Mutations of Perforin and Munc13-4 do not mark HLH by NK defects. *Pediatr. Blood Cancer* 46: 409-411.

## CHROMOSOMAL LOCATION

Genetic locus: Munc13d (mouse) mapping to 11 E2.

## PRODUCT

Munc13-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Munc13-4 shRNA Plasmid (m): sc-62652-SH and Munc13-4 shRNA (m) Lentiviral Particles: sc-62652-V as alternate gene silencing products.

For independent verification of Munc13-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62652A, sc-62652B and sc-62652C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Munc13-4 siRNA (m) is recommended for the inhibition of Munc13-4 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Munc13-4 gene expression knockdown using RT-PCR Primer: Munc13-4 (m)-PR: sc-62652-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.