



OATP-D siRNA (h): sc-62713

BACKGROUND

The organic anion transporter family of proteins mediate hepatic uptake of cardiac glycosides. OATP-D (organic anion transporter D), also known as SLC01B3 (solute carrier organic anion transporter family member 1B3), SLC21A11 (solute carrier family 21 member 11) or PGE1 transporter, is a 710 amino acid member of the organic anion transporter protein family. As a multi-pass membrane protein, OATP-D mediates the Na⁺-independent transport of vasopressin, prostaglandins (PG) E1 and E2, thyroxine (T4), deltorphin II and other organic anions, but not estrone-3-sulfate, DPDPE, taurocholate, DHEAS or digoxin. OATP-D is ubiquitously expressed with highest levels present in leukocytes and spleen. OATP-D is expressed as four isoforms produced by alternative splicing events.

REFERENCES

1. Hsiang, B., et al. 1999. A novel human hepatic organic anion transporting polypeptide (OATP2). Identification of a liver-specific human organic anion transporting polypeptide and identification of rat and human hydroxymethylglutaryl-CoA reductase inhibitor transporters. *J. Biol. Chem.* 274: 37161-37168.
2. Konig, J., et al. 2000. Localization and genomic organization of a new hepatocellular organic anion transporting polypeptide. *J. Biol. Chem.* 275: 23161-23168.
3. Cai, S.Y., et al. 2002. An evolutionarily ancient Oatp: insights into conserved functional domains of these proteins. *Am. J. Physiol. Gastrointest. Liver Physiol.* 282: G702-G710.
4. Pizzagalli, F., et al. 2002. Identification of a novel human organic anion transporting polypeptide as a high affinity thyroxine transporter. *Mol. Endocrinol.* 16: 2283-2296.
5. Patel, P., et al. 2002. Semi quantitative expression analysis of Mdr-3, FIC1, BSEP, OATP-A, OATP-C, OATP-D, OATP-E and NTCP gene transcripts in 1st and 3rd trimester human placenta. *Placenta* 24: 39-44.
6. Adachi, H., et al. 2003. Molecular characterization of human and rat organic anion transporter OATP-D. *Am. J. Physiol. Renal Physiol.* 285: F1188-F1197.
7. Niemi, M., et al. 2004. High plasma pravastatin concentrations are associated with single nucleotide polymorphisms and haplotypes of organic anion transporting polypeptide-C (Oatp-C, SLC01B1). *Pharmacogenetics* 14: 429-440.
8. SWISS-PROT/TrEMBL (Q9UIG8). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: SLC03A1 (human) mapping to 15q26.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

OATP-D siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see OATP-D shRNA Plasmid (h): sc-62713-SH and OATP-D shRNA (h) Lentiviral Particles: sc-62713-V as alternate gene silencing products.

For independent verification of OATP-D (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62713A, sc-62713B and sc-62713C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

OATP-D siRNA (h) is recommended for the inhibition of OATP-D expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor OATP-D gene expression knockdown using RT-PCR Primer: OATP-D (h)-PR: sc-62713-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.